



Absolutely convex sets of large Szlenk index

Philip A. H. Brooker

Received: October 21, 2018/Accepted: June 18, 2019/Online: August 16, 2019

Abstract

Let X be a Banach space and K an absolutely convex, weak*-compact subset of X^* . We study consequences of K having a large or undefined Szlenk index and subsequently derive a number of related results concerning basic sequences and universal operators. We show that if X has a countable Szlenk index then X admits a subspace Y such that Y has a basis and the Szlenk indices of Y are comparable to the Szlenk indices of X . If X is separable, then X also admits subspace Z such that the quotient X/Z has a basis and the Szlenk indices of X/Z are comparable to the Szlenk indices of X . We also show that for a given ordinal ξ the class of operators whose Szlenk index is not an ordinal less than or equal to ξ admits a universal element if and only if $\xi < \omega_1$; W.B. Johnson's theorem that the formal identity map from ℓ_1 to ℓ_∞ is a universal non-compact operator is then obtained as a corollary. Stronger results are obtained for operators having separable codomain.

Keywords: Szlenk index, Asplund space, basic sequence, universal operator.

msc: 46B20, 46B03, 46B15, 47B10.

1 Introduction

The Szlenk index is an ordinal index that measures the difference between the norm and weak* topologies on subsets of dual Banach spaces. It was introduced by Szlenk¹ to solve (in the negative) the problem of whether there exists a separable, reflexive Banach space whose subspaces exhaust the class of separable, reflexive Banach spaces up to isomorphism. Since then the Szlenk index has found many uses in the study of Banach spaces and their operators, as outlined in the surveys of Lancien (2006) and Rosenthal (2003).

In the current paper we study the Szlenk index in two main contexts, the first of these being the theory of basic sequences in Banach spaces. Our work on basic sequences and the Szlenk index is based on the classical method of Mazur² for producing subspaces with a basis and on the more recent method of Johnson

¹Szlenk, 1968, "The non-existence of a separable reflexive Banach space universal for all separable reflexive Banach spaces".

²See e.g. Albiac and Kalton, 2016, *Topics in Banach space theory*, Section 1.5.

and Rosenthal³ for producing quotients with a basis (a dual version of the Mazur technique). We extend previous work in this area by Lancien⁴ and Dilworth-Kutzarova-Lancien-Randrianarivony⁵. The other main motivation for our paper is the study of *universal* operators. In Sections 1.1 and 1.2 we provide relevant background on these two areas of study and explain the main contributions of the current paper to these topics, along the way providing an outline of the content of Sections 3 to 7. Basic notation and terminology for Banach space theory is set out in Section 1.3, whilst background information on more specialised topics such as Szlenk indices, trees and operators acting on Banach spaces over trees will be provided in Section 2.

1.1 Structure of Banach spaces with large Szlenk index

Several authors have already studied the structure of subspaces and quotients of Banach spaces having Szlenk index larger than a given ordinal. We shall now sketch the previous such results of most relevance to us and explain the contributions of the current paper to this topic. The first result of interest to us is the following result due to G. Lancien⁶.

Proposition 1 – *Let X be a Banach space and $\xi < \omega_1$. If $Sz(X) > \xi$ then there exists a separable subspace Y of X such that $Sz(Y) > \xi$.*

To prove Proposition 1 Lancien showed⁷ that for a suitable tree \mathcal{T} of rank $\xi + 1$, the estimate $Sz(X, \epsilon) > \xi$ (which follows for some $\epsilon > 0$ from the estimate $Sz(X) > \xi$) implies the existence of families of vectors $(x_t)_{t \in \mathcal{T}} \subseteq B_X$ and $(x_t^*)_{t \in \mathcal{T}} \subseteq B_{X^*}$ whose construction depends on ϵ and which satisfy certain properties that bear witness to the fact that $Sz(X) > \xi$. Without giving the precise details of Lancien’s construction here, we mention that the subspace Y is taken to be the closed linear span in X of the family $(x_t)_{t \in \mathcal{T}}$. Recently, Dilworth, Kutzarova, Lancien and Randrianarivony⁸ have adapted Lancien’s construction to show that, under the additional hypothesis that X is reflexive, $(x_t)_{t \in \mathcal{T}}$ may be assumed to be a basic sequence for a suitable enumeration of \mathcal{T} . In Section 3 we devise a further refinement of Lancien’s construction, the precise details of which are captured in the statement of our main technical result, Theorem 9. A number of consequences of Theorem 9 follow, including the main results of the current paper, namely Theorems 1, 2, 6 and 7.

Our first application of Theorem 9 is the following result, proved in Section 4, which asserts that if X is an infinite-dimensional Banach space with countable

³Johnson and Rosenthal, 1972, “On w^* -basic sequences and their applications to the study of Banach spaces”.

⁴Lancien, 1996, “On the Szlenk index and the weak*-dentability index”.

⁵Dilworth et al., 2017, “Equivalent norms with the property (β) of Rolewicz”.

⁶Lancien, 1996, “On the Szlenk index and the weak*-dentability index”, Proposition 3.1.

⁷Ibid., Lemma 3.4.

⁸Dilworth et al., 2017, “Equivalent norms with the property (β) of Rolewicz”, Proposition 3.1(i).

1. Introduction

Szlenk index, then X admits a subspace with a basis and with ϵ -Szlenk indices comparable to the ϵ -Szlenk indices of X :

Theorem 1 – *Let X be an infinite-dimensional Banach space such that $Sz(X) < \omega_1$ and let $\delta > 0$. Then there exists a subspace $Y \subseteq X$ such that Y has a shrinking basis with basis constant not exceeding $1 + \delta$ and such that*

$$\forall \epsilon > 0 \quad Sz\left(Y, \frac{\epsilon}{66}\right) \geq Sz(X, \epsilon), \quad (1)$$

hence $Sz(Y) = Sz(X)$.

To compare Theorem 1 with the earlier results of Lancien (1996) and Dilworth et al. (2017), first note that it follows easily from Proposition 1 that if X is a Banach space with $Sz(X) < \omega_1$, then there exists a separable subspace Y of X such that $Sz(Y) = Sz(X)$. For instance, take Y to be the closed linear span of $\bigcup_{n=1}^{\infty} Y_n$ in X , where, for each $n \in \mathbb{N}$, Y_n is a separable subspace of X with $Sz(Y_n) > Sz(X, 1/n) - 1$. However, even in the case that X is reflexive one cannot in general conclude from Proposition 3.1(i) of Dilworth et al. (2017) (or from Proposition 1) that X admits a subspace with a basis *and* with Szlenk index equal to the Szlenk index of X . Indeed, from Proposition 3.1(i) of Dilworth et al. (2017) we may only deduce that the subspace Y with a basis satisfies $Sz(Y) \geq \xi \omega$ (on the other hand, if X is reflexive and $Sz(X) = \omega^{\alpha+1}$ for some $\alpha < \omega_1$, then it follows from Proposition 3.1(i) of Dilworth et al. (2017) that X admits a subspace Y with a basis and satisfying $Sz(Y) = Sz(X)$; to see this, take a subspace Y with a basis such that $Sz(Y) > \omega^\alpha$).

One of the most important features of Theorem 9 is that the construction of families of vectors (x_t) and (x_t^*) in the proof of the theorem depends on a family $(\epsilon_n)_{n < \omega}$ of (possibly infinitely many distinct) epsilons and associated estimates $Sz(K, \epsilon_n) > \xi_n$; the earlier such constructions from Lancien (1996) and Dilworth et al. (2017) depend instead on consideration of just a single ϵ and an associated estimate $Sz(X, \epsilon) > \xi$. The greater generality taken in our approach plays a key role in the proof of Theorem 1. In particular whereas the constructions underpinning Proposition 1 above and Proposition 3.1(i) of Dilworth et al. (2017) yield an estimate of the type shown at Equation (1) for single $\epsilon > 0$, with our approach we take the sequence $(\epsilon_n)_{n < \omega}$ in the statement of Theorem 9 to be dense in $(0, \infty)$ and apply a straightforward density argument to obtain the estimate shown at Equation (1) for all $\epsilon > 0$, which is enough to conclude the final assertion of Theorem 1 (without any reflexivity hypothesis).

We now turn our attention to quotients. Under the additional assumption that X is separable, Theorem 9 yields the existence of a quotient of X having a basis and with ϵ -Szlenk indices comparable to the ϵ -Szlenk indices of X , as per the following result established in Section 4.

Theorem 2 – *Let X be an infinite-dimensional Banach space with separable dual and let $\delta > 0$. Then there exists a subspace $Z \subseteq X$ such that X/Z has a shrinking basis with*

basis constant not exceeding $1 + \delta$ and such that

$$\forall \epsilon > 0 \quad \text{Sz}\left(X/Z, \frac{\epsilon}{9}\right) \geq \text{Sz}(X, \epsilon), \tag{2}$$

hence $\text{Sz}(X/Z) = \text{Sz}(X)$.

Theorem 2 should be compared with the following earlier result of G. Lancien⁹.

Proposition 2 – *Let X be a separable Banach space and $\xi < \omega_1$. If $\text{Sz}(X) > \xi$ then there exists a subspace Z of X such that X/Z has a shrinking basis and $\text{Sz}(X/Z) > \xi$.*

To prove Proposition 2 Lancien combined the construction developed in the proof of Lemma 3.4 of Lancien (1996) (previously mentioned in the discussion following Proposition 1 on p. 164) with the techniques developed by Johnson and Rosenthal¹⁰ for constructing weak*-basic sequences in dual Banach spaces. We take a similar approach in the proof of Theorem 9, with the main difference between Theorem 2 and Proposition 2 again arising from the more general approach taken in the construction of the families of vectors (x_t) and (x_t^*) in the proof of the Theorem 9, in particular the dependence on an infinite family $(\epsilon_n)_{n < \omega}$ of (possibly distinct) epsilons and associated estimates $\text{Sz}(K, \epsilon_n) > \xi_n$ rather than just a single ϵ and associated estimate $\text{Sz}(X, \epsilon)$.

Our comparison of Theorem 2 with Proposition 2 is similar to our comparison of Theorem 1 with Proposition 1. In general one may deduce from Proposition 2 only that $\text{Sz}(X/Z) \geq \xi_\omega$; the stronger assertion that $\text{Sz}(X/Z) = \text{Sz}(X)$ need not follow except in the case that $\text{Sz}(X) = \omega^{\alpha+1}$ for some ordinal α and we take Z to satisfy $\text{Sz}(X/Z) > \omega^\alpha$. By contrast, Theorem 2 provides the conclusion that $\text{Sz}(X/Z) = \text{Sz}(X)$ in all cases.

Remark 1 – Trees have been used by a number of authors to study and compare a variety of ordinal indices on Banach spaces. For instance, the work of Alspach, Judd, and Odell (2005) defines and studies a number of local ℓ_1 -indices and compares these indices with one another and with the Szlenk index.

1.2 Universal operators

The initial motivation for writing the current paper was a desire to study the Szlenk index in the context of the problem of finding universal elements for certain subclasses of the class \mathcal{L} of all (bounded, linear) operators between Banach spaces. For operators $T \in \mathcal{L}(X, Y)$ and $S \in \mathcal{L}(W, Z)$, where W, X, Y and Z are Banach spaces, we say that S *factors through* T (or, equivalently, that T *factors* S) if there

⁹Lancien, 1996, “On the Szlenk index and the weak*-dentability index”, Proposition 3.5.

¹⁰Johnson and Rosenthal, 1972, “On w^* -basic sequences and their applications to the study of Banach spaces”.

1. Introduction

exist $U \in \mathcal{L}(W, X)$ and $V \in \mathcal{L}(Y, Z)$ such that $VTU = S$. With this terminology, for a given subclass \mathcal{C} of \mathcal{L} we say that an operator $\Upsilon \in \mathcal{C}$ is *universal for \mathcal{C}* if Υ factors through every element of \mathcal{C} . Typically \mathcal{C} will be the complement $\complement \mathcal{I}$ of an operator ideal \mathcal{I} in the sense of Pietsch¹¹ (that is, $\complement \mathcal{I}$ consists of all elements of \mathcal{L} that do not belong to \mathcal{I}), or perhaps the restriction $\mathcal{I} \cap \complement \mathcal{I}$ of $\complement \mathcal{I}$ to a large subclass \mathcal{J} of \mathcal{L} ; e.g., \mathcal{J} might denote a large operator ideal or the class of all operators having a specified domain or codomain. One may think of a universal element of the class \mathcal{C} as a minimal element of \mathcal{C} that is ‘fixed’ or ‘preserved’ by each element of \mathcal{C} .

The notion of universality for a class of operators goes back to the work of Lindenstrauss and Pełczyński¹², who obtained the following result.

Theorem 3 – *Let X and Y be Banach spaces and suppose $T : X \rightarrow Y$ is a non-weakly compact operator. Then T factors the (non-weakly compact) summation operator $\Sigma : (a_n)_{n=1}^\infty \mapsto (\sum_{i=1}^n a_i)_{n=1}^\infty$ from ℓ_1 to ℓ_∞ . In particular, Σ is universal for the class of non-weakly compact operators.*

Since the publication of Lindenstrauss and Pełczyński (1968) a number of results in a similar spirit to Theorem 3 have appeared in the literature. Perhaps the most well-known is the following result of W.B. Johnson¹³, which is a special case of Theorem 6 of the current paper.

Theorem 4 – *Let X and Y be Banach spaces and suppose $T : X \rightarrow Y$ is a non-compact operator. Then T factors the (non-compact) formal identity operator from ℓ_1 to ℓ_∞ . In particular, the formal identity operator from ℓ_1 to ℓ_∞ is universal for the class of non-compact operators.*

Another universality result of note, due to C. Stegall¹⁴, is the existence of a universal non-Asplund operator. The Asplund operators have several equivalent definitions in the literature; in the current paper we say that an operator $T : X \rightarrow Y$ is *Asplund* if $(T|_Z)^*(Y^*)$ is separable for any separable subspace $Z \subseteq X$. We refer the reader to Stegall (1981) for further properties and characterisations of Asplund operators. Stegall’s universal operator is defined in terms of the Haar system $(h_m)_{m=0}^\infty \subseteq C(\{0, 1\}^\omega)$, where each factor $\{0, 1\}$ is discrete and $\{0, 1\}^\omega$ is equipped with its compact Hausdorff product topology. For the purpose of stating Stegall’s result, we let μ denote the product measure on $\{0, 1\}^\omega$ obtained by equipping each factor $\{0, 1\}$ with its discrete uniform probability measure and let $H : \ell_1 \rightarrow L_\infty(\{0, 1\}^\omega, \mu)$ be defined by setting $Hx = \sum_{m=1}^\infty x(m)h_{m-1}$ for each $x = (x(m))_{m=1}^\infty \in \ell_1$. Stegall’s result is the following theorem.

¹¹Pietsch, 1980, *Operator ideals*.

¹²Lindenstrauss and Pełczyński, 1968, “Absolutely summing operators in \mathcal{L}_p -spaces and their applications”, Theorem 8.1.

¹³Johnson, 1971, “A universal non-compact operator”.

¹⁴Stegall, 1975, “The Radon-Nikodým property in conjugate Banach spaces”, Theorem 4.

Theorem 5 – *Let X and Y be Banach spaces such that X is separable and suppose $T \in \mathcal{L}(X, Y)$ is such that $T^*(Y^*)$ is nonseparable. Then H factors through T .*

Since the domain of H , namely ℓ_1 , is separable, and since H^* has non-separable range, H is a non-Asplund operator. It therefore follows from Theorem 5 that H is a universal non-Asplund operator. We note that a different universal non-Asplund operator has been obtained by the author of the current paper using the techniques developed here in the context of studying the Szlenk index¹⁵.

Other universality results besides those mentioned above can be found in the work of Brooker (2017), Cilia and Gutiérrez (2015), Dilworth (1985), Girardi and Johnson (1997), Hinrichs and Pietsch (2000), Oikhberg (2016), and the Handbook survey on operator ideals by Diestel, Jarchow, and Pietsch (2001). Theorem 4 above has been applied in the study of information-based complexity by Hinrichs, Novak, and Woźniakowski (2013).

Remark 2 – K. Beanland and R.M. Causey¹⁶ have recently attained universality results for further classes of operators defined in terms of operator ideals.

In Section 5 we turn our attention to applying Theorem 9 to the study of universal operators for classes defined in terms of a bound on the Szlenk index. Our main result to this end is the following theorem.

Theorem 6 – *Let X and Y be Banach spaces, $\alpha < \omega_1$, $T \in \mathcal{L}(X, Y) \setminus \mathcal{SZ}_\alpha(X, Y)$ and (\mathcal{T}, \preceq) a countably infinite, rooted, well-founded tree with $\rho(\mathcal{T}) < \omega^{\alpha+1}$. Then $\Sigma_{\mathcal{T}^*}$ factors through T . Moreover if \mathcal{T} is blossomed and $\rho(\mathcal{T}) \geq \omega^\alpha$, then $\Sigma_{\mathcal{T}^*}$ is universal for \mathcal{CSZ}_α . It follows that, for an ordinal β , the class \mathcal{CSZ}_β admits a universal element if and only if $\beta < \omega_1$.*

In the statement of Theorem 6 \mathcal{SZ}_α denotes the class consisting of all operators whose Szlenk index is an ordinal no larger than ω^α (c.f. Definition 1 on p. 172), whilst the operator $\Sigma_{\mathcal{T}^*}$ is the path-sum operator of \mathcal{T}^* defined in Section 2.3, where \mathcal{T}^* is the rooted tree \mathcal{T} minus its (unique) minimal element.

Stronger universality results are achieved in Section 6 for operators having separable codomain, the main result in this direction being the following theorem.

Theorem 7 – *Let X and Y be Banach spaces, $\alpha < \omega_1$, $T \in \mathcal{L}(X, Y) \setminus \mathcal{SZ}_\alpha(X, Y)$ and (\mathcal{T}, \preceq) a countably infinite, rooted, well-founded tree with $\rho(\mathcal{T}) < \omega^{\alpha+1}$. If Y is separable then $\mathring{\sigma}_{\mathcal{T}}$ factors through T . Moreover if \mathcal{T} is blossomed and $\rho(\mathcal{T}) \geq \omega^\alpha$ then $\mathring{\sigma}_{\mathcal{T}}$ is universal for the class of non- α -Szlenk operators having separable codomain.*

The operator $\mathring{\sigma}_{\mathcal{T}}$ featured in the statement of Theorem 7 is the path-sum operator defined at Definition 4 on p. 198, which is essentially the same as $\Sigma_{\mathcal{T}}$ but with

¹⁵Brooker, 2017, “Non-Asplund Banach spaces and operators”.

¹⁶Beanland and Causey, 2017, “Genericity and universality for operator ideals”, arXiv:1711.09244.

1. Introduction

a restricted (in particular, norm separable) codomain. We conclude Section 6 by noting that stronger versions of the classical universality results of Lindenstrauss and Pełczyński (Theorem 3 on p. 167) and Johnson (Theorem 4 on p. 167) hold for operators whose codomain has weak*-sequentially compact dual ball.

In Section 7 we investigate whether the techniques developed in Sections 3 and 5 can be used to show the existence of universal operators for classes of Asplund operators having an uncountable strict lower bound for the Szlenk index. Though we do not completely answer this question, we show that the techniques developed in the earlier sections of the paper cannot decide the existence of universal operators in this setting in ZFC.

1.3 Notation and terminology

We now outline the notation and terminology used in the current paper. We work with Banach spaces over the scalar field $\mathbb{K} = \mathbb{R}$ or \mathbb{C} . Typical Banach spaces are denoted by the letters W, X, Y and Z , with the identity operator of X denoted Id_X . Following the terminology introduced earlier in the current paper, for Banach spaces W, X and Z and an operator $S : W \rightarrow Z$, we say that S *factors through* X if S factors through Id_X . We write X^* for the dual space of X and denote by ι_X the canonical embedding of X into X^{**} . We define $B_X := \{x \in X \mid \|x\| \leq 1\}$ and $B_X^\circ := \{x \in X \mid \|x\| < 1\}$. By a *subspace* of a Banach space X we mean a linear subspace of X that is closed in the norm topology. For a Banach space X , $C \subseteq X$ and $D \subseteq X^*$ we define $C^\perp := \{x^* \in X^* \mid \forall x \in C, x^*(x) = 0\}$ and $D_\perp = \{x \in X \mid \forall x^* \in D, x^*(x) = 0\}$. We denote by $[C]$ the norm closed linear hull of C in X , with a typical variation on this notation being that for an indexed set $\{x_i \mid i \in I\} \subseteq X$ we may write $[x_i]_{i \in I}$ or $[x_i \mid i \in I]$ in place of $[\{x_i \mid i \in I\}]$. With this notation we have that $\overline{[D]}^{w^*}$ is the weak* closed linear hull of D in X^* . We shall make use of the well-known fact that, for a Banach space X and a sequence $(x_m^*)_{m=1}^\infty \subseteq X^*$, the quotient map $Q : X \rightarrow X / \bigcap_{m=1}^\infty \ker(x_m^*)$ has the property that Q^* is an isometric weak*-isomorphism of $(X / \bigcap_{m=1}^\infty \ker(x_m^*))^*$ onto $\overline{[x_m^* \mid m \in \mathbb{N}]}^{w^*}$.

Operator ideals are denoted by script letters such as \mathcal{F} . Operator ideals of particular interest in the current paper are:

- \mathcal{K} , the *compact* operators;
- \mathcal{W} , the *weakly compact* operators;
- \mathcal{R} , the operators having separable range;
- \mathcal{R}^* , the operators whose adjoint has separable range;
- \mathcal{S} , the strictly singular operators;

- \mathcal{D} , the Asplund operators (also known as the *decomposing* operators); and,
- \mathcal{SZ}_α , the α -Szlenk operators for a given ordinal α .

All of the operator ideals in the list above are closed, and most of them are well known. An operator $T : X \rightarrow Y$ is *strictly singular* if for every infinite dimensional subspace $Z \subseteq X$ the restriction $T|_Z$ fails to be an isomorphic embedding of Z into Y (that is, if for every such Z and every $\epsilon > 0$ there exists $z \in Z$ such that $\|z\| = 1$ and $\|Tz\| < \epsilon$). For a given ordinal α , the class \mathcal{SZ}_α consists of all operators whose Szlenk index is an ordinal not exceeding ω^α . These classes have been studied in some detail by the current author¹⁷ and we note in particular that important relationships between the operator ideals \mathcal{SZ}_α and other ideals in the list above shall be given in Section 2.1. It is well known¹⁸ that \mathcal{X}^* is a subclass of \mathcal{X} .

For infinite dimensional Banach spaces X and Z we say that X is *Z-saturated* if every infinite dimensional subspace of X contains a further subspace that is isomorphic to Z . It is well known¹⁹ that for each $1 \leq p < \infty$ the Banach space ℓ_p is ℓ_p -saturated and that for distinct $p, q \in [1, \infty)$ neither of the spaces ℓ_p and ℓ_q is isomorphic to a subspace of the other. From this it follows readily that if X is ℓ_p -saturated and Y is ℓ_q -saturated for distinct $p, q \in [1, \infty)$ then every operator from X to Y is strictly singular (we shall use this fact later in the proof of Proposition 10).

By **Ord** we denote the class of all ordinals, so that by $\alpha \in \mathbf{Ord}$ we mean that α is an ordinal. We write $\text{cof}(\alpha)$ for the cofinality of the ordinal α . If α is a successor ordinal, we write $\alpha - 1$ to mean the unique ordinal whose successor is α .

For a set S and a subset $R \subseteq S$ we write χ_R^S for the indicator function of R in S , or simply χ_R if no confusion can result. When discussing a Banach space $\ell_1(S)$ for some set S , for $s \in S$ we typically denote by e_s the element of $\ell_1(S)$ satisfying $e_s(s') = 1$ if $s' = s$ and $e_s(s') = 0$ if $s' \neq s$ ($s' \in S$). We thus denote by $(e_n)_{n=1}^\infty$ the standard unit vector basis of $\ell_1 = \ell_1(\mathbb{N})$. Where confusion may otherwise result, we may write e_s^S in place of e_s to specify the space $\ell_1(S)$ to which e_s belongs.

We shall repeatedly use the fact that for a set I , Banach space X and family $\{x_i \mid i \in I\} \subseteq X$ with $\sup_{i \in I} \|x_i\| < \infty$, there exists a unique element of $\mathcal{L}(\ell_1(I), X)$ satisfying $e_i \mapsto x_i$, $i \in I$.

For a Banach space X , a subset $A \subseteq X$, and $\epsilon > 0$, we say that A is ϵ -separated if $\|x - y\| > \epsilon$ for any distinct $x, y \in A$. For $B \subseteq C \subseteq X$ and $\delta > 0$ we say that B is a δ -net in C if for every $w \in C$ there exists $z \in B$ such that $\|w - z\| \leq \delta$.

¹⁷Brooker, 2012, "Asplund operators and the Szlenk index".

¹⁸See e.g. Pietsch, 1980, *Operator ideals*, Proposition 4.4.8.

¹⁹See e.g. Albiac and Kalton, 2016, *Topics in Banach space theory*, Chapter 2.

2 The Szlenk index, trees and operators on Banach spaces over trees

2.1 The Szlenk index

Let X be a Banach space. For each $\epsilon > 0$ define a derivation s_ϵ on weak*-compact subsets of X^* as follows: for weak*-compact $K \subseteq X^*$ let

$$s_\epsilon(K) := \{x^* \in K \mid \text{diam}(\mathcal{U} \cap K) > \epsilon \text{ for every weak}^*\text{-open } \mathcal{U} \ni x^*\}.$$

Iterate s_ϵ transfinitely by setting $s_\epsilon^0(K) = K$, $s_\epsilon^{\xi+1}(K) = s_\epsilon(s_\epsilon^\xi(K))$ for every ordinal ξ , and $s_\epsilon^\xi(K) = \bigcap_{\zeta < \xi} s_\epsilon^\zeta(K)$ whenever ξ is a limit ordinal. The ϵ -Szlenk index of K , denoted $\text{Sz}(K, \epsilon)$, is defined as the smallest ordinal ξ such that $s_\epsilon^\xi(K) = \emptyset$, if such an ordinal exists; if no such ordinal exists then $\text{Sz}(K, \epsilon)$ is undefined. (Note that, by weak*-compactness, $\text{Sz}(K, \epsilon)$ is a successor ordinal when it exists.) Notationally, we write $\text{Sz}(K, \epsilon) < \infty$ to mean that $\text{Sz}(K, \epsilon)$ is defined, and $\text{Sz}(K, \epsilon) = \infty$ to mean that $\text{Sz}(K, \epsilon)$ is undefined. If $\text{Sz}(K, \epsilon)$ is defined for all $\epsilon > 0$ then the Szlenk index of K , denoted $\text{Sz}(K)$, is the ordinal $\sup_{\epsilon > 0} \text{Sz}(K, \epsilon)$. If $\text{Sz}(K, \epsilon)$ is undefined for some $\epsilon > 0$, then $\text{Sz}(K)$ is undefined; we write $\text{Sz}(K) < \infty$ to mean that $\text{Sz}(K)$ is defined, and $\text{Sz}(K) = \infty$ to mean that $\text{Sz}(K)$ is undefined. Note that while $\text{Sz}(K, \epsilon) \leq \xi$ means that $\text{Sz}(K, \epsilon)$ is defined and equal to an ordinal not exceeding ξ , the statement $\text{Sz}(K, \epsilon) \not\leq \xi$ means either that $\text{Sz}(K, \epsilon)$ is undefined or that $\text{Sz}(K, \epsilon)$ is defined and exceeds ξ ; similarly, $\text{Sz}(K) \not\leq \xi$ means either that $\text{Sz}(K)$ is undefined or that $\text{Sz}(K)$ is defined and equal to an ordinal exceeding ξ .

Define the ϵ -Szlenk index of X and the Szlenk index of X to be the indices $\text{Sz}(X, \epsilon) := \text{Sz}(B_{X^*}, \epsilon)$ and $\text{Sz}(X) := \text{Sz}(B_{X^*})$, respectively. If Y is a Banach space and $T : X \rightarrow Y$ an operator, define the ϵ -Szlenk index of T and the Szlenk index of T to be the indices $\text{Sz}(T, \epsilon) := \text{Sz}(T^*(B_{X^*}), \epsilon)$ and $\text{Sz}(T) := \text{Sz}(T^*(B_{X^*}))$, respectively.

A useful survey of the Szlenk index and its applications in Banach space theory has been written by Lancien (2006). For facts regarding Szlenk indices of operators we refer the reader to the work of Brooker (2012). The following proposition collects some well-known facts concerning Szlenk indices of Banach spaces and operators.

Proposition 3 – *Let X be a Banach space.*

- (i) For $K_1 \subseteq K_2 \subseteq X^*$, $\epsilon_1 \geq \epsilon_2 > 0$ and ordinals $\xi_1 \geq \xi_2$ we have $s_{\epsilon_1}^{\xi_1}(K_1) \subseteq s_{\epsilon_2}^{\xi_2}(K_2)$.
- (ii) For a subspace $Z \subseteq X$ we have $\text{Sz}(Z) \leq \text{Sz}(X)$ and $\text{Sz}(X/Z) \leq \text{Sz}(X)$.
- (iii) The following are equivalent:
 - (a) $\text{Sz}(X) < \infty$ (that is, the Szlenk index is defined).
 - (b) X is an Asplund space.

- (c) X^* has the Radon-Nikodým property.
- (d) Every separable subspace of X has separable dual.

An argument due to A. Sersouri²⁰ shows that if the Szlenk index of a Banach space or an operator is defined, then it is of the form ω^α for some ordinal α (and more recently Causey²¹ has described all the possible ordinals α for which ω^α may be realised as the Szlenk index of a Banach space); this observation leads to the following definition.

Definition 1 – For α an ordinal define the class

$$\mathcal{SZ}_\alpha := \{T \in \mathcal{L} \mid \text{Sz}(T) \leq \omega^\alpha\}.$$

If $T \in \mathcal{SZ}_\alpha$, we say that T is α -Szlenk.

It is a result of Brooker (2012) that the classes \mathcal{SZ}_α are distinct for different values of α and that each such class is a closed operator ideal. Moreover, \mathcal{SZ}_0 coincides with the class \mathcal{K} of compact operators, whilst the class $\bigcup_{\alpha \in \text{Ord}} \mathcal{SZ}_\alpha$ of all operators whose Szlenk index is defined coincides with the class \mathcal{D} of Asplund operators. For operators with separable range, the following result of Brooker²² provides information regarding the relationship between the classes \mathcal{X}^* , \mathcal{D} and \mathcal{SZ}_α for $\alpha \in \text{Ord}$.

Proposition 4 – The following chain of equalities holds:

$$\mathcal{X}^* = \mathcal{K} \cap \mathcal{D} = \mathcal{K} \cap \bigcup_{\alpha \in \text{Ord}} \mathcal{SZ}_\alpha = \mathcal{K} \cap \bigcup_{\alpha < \omega_1} \mathcal{SZ}_\alpha = \mathcal{K} \cap \mathcal{SZ}_{\omega_1}.$$

2.2 Trees

A *tree* is a partially ordered set (T, \leq) for which the set $\{s \in T \mid s \leq t\}$ is well-ordered for every $t \in T$. We shall frequently suppress the partial order \leq and refer to the underlying set T as the tree. An element of a tree is called a *node*. For $S \subseteq T$ we denote by $\text{MIN}(S)$ (resp., $\text{MAX}(S)$) the set of all minimal (resp., maximal) elements of S . A *subtree* of T is a subset of T equipped with the partial order induced by the partial order of T , which we also denote \leq . A *chain* in T is a totally ordered subset of T . A *branch* of T is a maximal (with respect to set inclusion) totally ordered subset of T . We say that T is *well-founded* if it contains no infinite branches, and *chain-complete* if every chain \mathcal{C} in T admits a unique least upper bound. Clearly, every well-founded tree is chain-complete. A subset $S \subseteq T$ is said to be *downwards closed* in T if $S = \bigcup_{t \in S} \{s \in T \mid s \leq t\}$. Following Todorčević (1984), a *path* in T is a downwards

²⁰Sersouri, 1989, “Lavrientiev index for Banach spaces”, proof of Lemma 6.

²¹Causey, 2017, “The Szlenk index of injective tensor products and convex hulls”.

²²Brooker, 2012, “Asplund operators and the Szlenk index”, Proposition 2.11.

2. The Szlenk index, trees and operators on Banach spaces over trees

closed, totally ordered subset of \mathcal{T} . An *interval* in \mathcal{T} is a subset of \mathcal{T} of the form $(t', t'']$, $[t', t'']$, $[t', t'')$ or (t', t'') , where, for $t', t'' \in \mathcal{T}$, $(t', t'') := \{t \in \mathcal{T} \mid t' < t \leq t''\}$ and the other types of intervals are defined analogously. (For a tree (\mathcal{T}, \leq) and $s, t \in \mathcal{T}$ we write $s < t$ to mean that $s \leq t$ and $s \neq t$.) For $t \in \mathcal{T}$ we define the following sets:

$$\begin{aligned} \mathcal{T}[\leq t] &= \{s \in \mathcal{T} \mid s \leq t\} \\ \mathcal{T}[< t] &= \{s \in \mathcal{T} \mid s < t\} \\ \mathcal{T}[t \leq] &= \{s \in \mathcal{T} \mid t \leq s\} \\ \mathcal{T}[t <] &= \{s \in \mathcal{T} \mid t < s\} \\ \mathcal{T}[t+] &= \text{MIN}(\mathcal{T}[t <]) \end{aligned}$$

By t^- we denote the maximal element of $\mathcal{T}[< t]$, if it exists (that is, if the order type of $\mathcal{T}[< t]$ is a successor). If $s, t \in \mathcal{T}$ are such that $s \not\leq t$ and $t \not\leq s$, then we write $s \perp t$. Following Godefroy, Kalton, and Lancien (2001), a subtree \mathcal{S} of \mathcal{T} is said to be a *full* subtree of \mathcal{T} if it is downwards closed, $|\mathcal{S} \cap \text{MIN}(\mathcal{T})| = |\text{MIN}(\mathcal{T})|$, and for every $t \in \mathcal{S}$ we have $|\mathcal{S}[t+]| = |\mathcal{T}[t+]|$. A tree is said to be *rooted* if $|\text{MIN}(\mathcal{T})| \leq 1$. In particular, a nonempty tree is rooted if and only if it admits a unique minimal element, which we call the *root* of \mathcal{T} . We denote by \mathcal{T}^* the subtree $\mathcal{T} \setminus \text{MIN}(\mathcal{T})$ of \mathcal{T} . For $t \in \mathcal{T}$ the *height* of t , denoted $ht_{\mathcal{T}}(t)$, is the order type of $\mathcal{T}[< t]$. The *height* of \mathcal{T} is the ordinal $ht(\mathcal{T}) = \sup\{ht_{\mathcal{T}}(t) + 1 \mid t \in \mathcal{T}\}$. Note that $ht(\mathcal{T}) \leq \omega$ if and only if $\mathcal{T}[< t]$ is finite for every $t \in \mathcal{T}$.

Let $\mathcal{T} = (\mathcal{T}, \leq)$ be a tree, α an ordinal and $\psi : \alpha \rightarrow \mathcal{T}$ a surjection. Then ψ induces a well-ordering of \mathcal{T} that extends \leq . Indeed, define $A_0 = \mathcal{T}[\leq \psi(0)]$ and, if $\beta > 0$ is an ordinal such that A_γ has been defined for all $\gamma < \beta$, define $A_\beta = \mathcal{T}[\leq \psi(\beta)] \setminus \bigcup_{\gamma < \beta} \mathcal{T}[\leq \psi(\gamma)]$. The induced well-order \leq of \mathcal{T} is defined by declaring $s \leq t$, where $s \in A_\beta$ and $t \in A_{\beta'}$, if $\beta < \beta'$ or if $\beta = \beta'$ and $s \leq t$. Note that if \mathcal{T} is countable and $ht(\mathcal{T}) \leq \omega$ then the well-ordering of \mathcal{T} induced as above by a surjection of ω onto \mathcal{T} is of order type ω . In fact, the following statements are equivalent for an infinite tree \mathcal{T} :

- (i₀) \mathcal{T} is countable and $\mathcal{T}[< t]$ is finite for every $t \in \mathcal{T}$;
- (ii₀) \mathcal{T} is countable and $ht(\mathcal{T}) \leq \omega$;
- (iii₀) There exists a bijection τ of ω onto \mathcal{T} such that $\tau(l) \leq \tau(m)$ implies $l \leq m$ for $l, m < \omega$.

Example 1 – Let $\Omega := \bigcup_{n < \omega} \prod_n \omega$. That is, Ω is the set of all finite (including possibly empty) sequences of finite ordinals. We define an order \sqsubseteq on Ω by saying that $s \sqsubseteq t$ if and only if s is an initial segment of t . Note that Ω is a rooted tree, with its root being the empty sequence \emptyset . For $n < \omega$ and $t \in \Omega$ we denote by $n^\frown t$ the concatenation of (n) with t ; that is, $n^\frown t = (n)$ if $t = \emptyset$ and $n^\frown t = (n, n_1, \dots, n_k)$ if $t = (n_1, \dots, n_k)$. It is straightforward to show that for an arbitrary tree (\mathcal{T}, \leq) the following statements are equivalent to statements (i₀)–(iii₀) above and to each other:

(iv₀) \mathcal{T} is order-isomorphic to a subtree of Ω ;

(v₀) \mathcal{T} is order-isomorphic to a downwards-closed subtree of Ω^\star .

Moreover, if \mathcal{T} is rooted then (i₀)–(iv₀) are equivalent to:

(vi₀) \mathcal{T} is order-isomorphic to a downwards-closed subtree of Ω .

We now describe a method for inductively defining a decreasing (with respect to set inclusion) family of downwards closed subtrees of a given tree, indexed by the ordinals. To this end for a tree (\mathcal{T}, \leq) let

$$\begin{aligned} \mathcal{T}^{(0)} &= \mathcal{T}; \\ \mathcal{T}^{(\xi+1)} &= \mathcal{T}^{(\xi)} \setminus \text{MAX}(\mathcal{T}^{(\xi)}) \quad \text{for every ordinal } \xi; \text{ and,} \\ \mathcal{T}^{(\xi)} &= \bigcap_{\xi' < \xi} \mathcal{T}^{(\xi')} \quad \text{if } \xi \text{ is a limit ordinal.} \end{aligned}$$

The fact that $\mathcal{T}^{(\xi)}$ is downwards closed in \mathcal{T} for all ordinals ξ follows from a straightforward transfinite induction.

The *rank* of a node $t \in \mathcal{T}$ is defined to be the unique ordinal $\rho_{\mathcal{T}}(t)$ such that $t \in \mathcal{T}^{(\rho_{\mathcal{T}}(t))} \setminus \mathcal{T}^{(\rho_{\mathcal{T}}(t)+1)}$, if it exists. Notice that if $s, t \in \mathcal{T}$ are such that $t < s$ and $\rho_{\mathcal{T}}(t)$ exists, then $\rho_{\mathcal{T}}(s)$ exists and satisfies $\rho_{\mathcal{T}}(s) < \rho_{\mathcal{T}}(t)$ since the derived trees $\mathcal{T}^{(\xi)}$ are downwards closed. It follows that if $t \in \mathcal{T}$ is such that $\rho_{\mathcal{T}}(s)$ exists for all $s \in \mathcal{T}[t+]$, then $\rho_{\mathcal{T}}(t)$ exists and satisfies

$$\rho_{\mathcal{T}}(t) = \sup\{\rho_{\mathcal{T}}(s) + 1 \mid s \in \mathcal{T}[t+]\}. \tag{3}$$

Thus, if $t_0 \in \mathcal{T}$ is such that $\rho_{\mathcal{T}}(t_0)$ does not exist, there exists $t_1 \in \mathcal{T}[t_0+]$ such that $\rho_{\mathcal{T}}(t_1)$ does not exist; similarly, there exists $t_2 \in \mathcal{T}[t_1+]$ such that $\rho_{\mathcal{T}}(t_2)$ does not exist, and in this way we inductively define an infinite chain $(t_n)_{n < \omega}$ in \mathcal{T} , hence \mathcal{T} is not well-founded. Conversely, if \mathcal{T} contains an infinite branch, \mathcal{B} say, then, with s_n denoting the element of \mathcal{B} of height n in \mathcal{T} , we have by induction that $\{s_n \mid n < \omega\} \subseteq \mathcal{T}^{(\xi)}$ for all ordinals ξ , hence $\rho_{\mathcal{T}}(s_n)$ is undefined for all n . We deduce that $\rho_{\mathcal{T}}(t)$ exists for all $t \in \mathcal{T}$ if and only if \mathcal{T} is well-founded, if and only if $\mathcal{T}^{(\xi)} = \emptyset$ for some ordinal ξ . If \mathcal{T} is well-founded, the *rank* of \mathcal{T} is the ordinal

$$\rho(\mathcal{T}) := \min\{\xi \mid \mathcal{T}^{(\xi)} = \emptyset\} = \sup\{\rho_{\mathcal{T}}(t) + 1 \mid t \in \mathcal{T}\}.$$

Notice that if \mathcal{T} is rooted, with root t_0 , say, then $\mathcal{T}^{(\rho_{\mathcal{T}}(t_0))} = \{t_0\}$, hence $\rho(\mathcal{T}) = \rho_{\mathcal{T}}(t_0) + 1$ is a successor ordinal.

We now give the definition of a *blossomed* tree, due to Gasparis (2005).

Definition 2 – We say that a countable tree (\mathcal{T}, \leq) is *blossomed* if it is rooted, well-founded, and for each $t \in \mathcal{T} \setminus \text{MAX}(\mathcal{T})$ there exists a bijection $\psi : \omega \rightarrow \mathcal{T}[t+]$ such that $m \leq n < \omega$ implies $\rho_{\mathcal{T}}(\psi(m)) \leq \rho_{\mathcal{T}}(\psi(n))$.

2. The Szlenk index, trees and operators on Banach spaces over trees

Blossomed trees were used by Gasparis²³ to study fixing properties of operators of large Szlenk index acting on $C(K)$ spaces. The important property of a blossomed tree \mathcal{T} in studying the Szlenk index is that for every $t \in \mathcal{T} \setminus \text{MAX}(\mathcal{T})$ and cofinite subset $\mathcal{Q} \subseteq \mathcal{T}[t+]$ we have

$$\sup\{\rho_{\mathcal{T}}(t') \mid t' \in \mathcal{Q}\} = \sup\{\rho_{\mathcal{T}}(t'') \mid t'' \in \mathcal{T}[t+]\}; \quad (4)$$

this condition clearly holds for any blossomed tree and, moreover, any countable, rooted, well-founded tree \mathcal{T} satisfying the property stated at Equation (4) admits a subtree \mathcal{S} such that $\rho(\mathcal{S}) = \rho(\mathcal{T})$ and \mathcal{S} is blossomed. Thus, blossomed trees can be thought of as the ‘minimal’ trees satisfying these conditions and, moreover, the formally stronger definition of a blossomed tree is typically more convenient to work with than the property stated at Equation (4) for the purposes of proving results concerning the Szlenk index, such as Proposition 11.

The following example guarantees a rich supply of blossomed trees. Note that other examples of blossomed trees, namely the Schreier families of finite subsets of \mathbb{N} , are used in the context of Szlenk indices in Gasparis (2005). The construction of trees in Example 2 below is essentially the same as that given by Bourgain on p. 91 of Bourgain (1979).

Example 2 – We construct, via transfinite induction on $\xi < \omega_1$, a family $(\mathcal{T}_\xi)_{\xi < \omega_1}$ consisting of blossomed subtrees of Ω that satisfy $\rho(\mathcal{T}_\xi) = \xi + 1$ for each $\xi < \omega_1$. Set $\mathcal{T}_0 = \{\emptyset\}$. Suppose $\xi > 0$ is an ordinal such that the \mathcal{T}_ζ has been defined for all $\zeta < \xi$; we define \mathcal{T}_ξ as follows. Let $(\xi_n)_{n=0}^\infty$ be a non-decreasing, cofinal sequence in ξ , and set

$$\mathcal{T}_\xi = \{\emptyset\} \cup \{n \frown t \mid n < \omega, t \in \mathcal{T}_{\xi_n}\}.$$

A straightforward transfinite induction on $\zeta \leq \xi$ shows that

$$\forall \zeta \leq \xi \quad \mathcal{T}_\xi^{(\zeta)} = \{\emptyset\} \cup \bigcup_{n < \omega} \{n \frown t \mid t \in \mathcal{T}_{\xi_n}^{(\zeta)}\}, \quad (5)$$

hence

$$\forall \zeta < \xi \quad \text{MAX}(\mathcal{T}_\xi^{(\zeta)}) = \bigcup_{n < \omega} \{n \frown t \mid t \in \text{MAX}(\mathcal{T}_{\xi_n}^{(\zeta)})\}. \quad (6)$$

Taking $\zeta = \xi$ in Equation (5) yields $\mathcal{T}_\xi^{(\xi)} = \{\emptyset\}$, hence $\rho_{\mathcal{T}_\xi}(\emptyset) = \xi$ and $\rho(\mathcal{T}_\xi) = \xi + 1$.

For $n < \omega$ let $\iota_n : \mathcal{T}_{\xi_n} \rightarrow \mathcal{T}_\xi$ be the map $t \mapsto n \frown t$. From Equation (6) we have $\rho_{\mathcal{T}_\xi}(n \frown t) = \rho_{\mathcal{T}_{\xi_n}}(t)$ every $n < \omega$ and $t \in \mathcal{T}_{\xi_n}$. Thus, if for $n < \omega$ and $t \in \mathcal{T}_{\xi_n} \setminus \text{MAX}(\mathcal{T}_{\xi_n})$ the map $\psi : \omega \rightarrow \mathcal{T}_{\xi_n}[t+]$ is a bijection such that $(\rho_{\mathcal{T}_{\xi_n}}(\psi(m)))_{m=0}^\infty$ is non-decreasing (as per Example 2), then $\iota_n \circ \psi : \omega \rightarrow \mathcal{T}_\xi[(n \frown t)+]$ is a bijection with

$$(\rho_{\mathcal{T}_\xi}(\iota_n \circ \psi(m)))_{m=0}^\infty = (\rho_{\mathcal{T}_\xi}(n \frown \psi(m)))_{m=0}^\infty = (\rho_{\mathcal{T}_{\xi_n}}(\psi(m)))_{m=0}^\infty$$

²³Gasparis, 2005, “Operators on $C(K)$ spaces preserving copies of Schreier spaces”.

non-decreasing. Similarly, $n \mapsto (n)$ is a bijection of ω onto $\mathcal{T}_\xi[\emptyset+]$ and

$$(\rho_{\mathcal{T}_\xi}((n)))_{n=0}^\infty = (\rho_{\mathcal{T}_{\xi_n}}(\emptyset))_{n=0}^\infty = (\xi_n)_{n=0}^\infty$$

is non-decreasing. We have now shown that \mathcal{T}_ξ is blossomed, as required.

The following proposition collects properties of blossomed trees that we shall need in subsequent sections of the current paper.

Proposition 5 – *Let (\mathcal{S}, \leq') and (\mathcal{T}, \leq) be countable, rooted, well-founded trees.*

- (i) *If \mathcal{S} is blossomed and $\rho(\mathcal{T}) \leq \rho(\mathcal{S})$ then \mathcal{T} is order isomorphic to a downwards closed subtree of \mathcal{S} .*
- (ii) *If \mathcal{S} is blossomed and \mathcal{S}' is a full subtree of \mathcal{S} then \mathcal{S}' is blossomed and $\rho(\mathcal{S}') = \rho(\mathcal{S})$.*

Assertion (i) of Proposition 5 is a trivial generalisation of Lemma 2.7 of Gasparis (2005), requiring little change in the proof. Assertion (ii) is Lemma 2.8 of Gasparis (2005). We refer the reader to Gasparis (2005) for the proofs.

There are various natural topologies for trees, many of which are described in Nyikos (1997). The tree topology which will be of interest to us in Section 6 is the *coarse wedge topology*, which is compact and Hausdorff for many trees. The coarse wedge topology of (\mathcal{T}, \leq) is that topology on \mathcal{T} formed by taking as a subbase all sets of the form $\mathcal{T}[t \leq]$ and $\mathcal{T} \setminus \mathcal{T}[t \leq]$, where the order type of $\mathcal{T}[< t]$ is either 0 or a successor ordinal. For a tree (\mathcal{T}, \leq) , $t \in \mathcal{T}$ and $\mathcal{F} \subseteq \mathcal{T}$, define

$$\mathcal{W}_T(t, \mathcal{F}) := \mathcal{T}[t \leq] \setminus \bigcup_{s \in \mathcal{F}} \mathcal{T}[s \leq].$$

The following proposition is clear.

Proposition 6 – *Let (\mathcal{T}, \leq) be a tree and let $t \in \mathcal{T}$ be such that the order type of $\mathcal{T}[< t]$ is 0 or a successor ordinal. Then the coarse wedge topology of \mathcal{T} admits a local base of clopen sets at t consisting of all sets of the form $\mathcal{W}_T(t, \mathcal{F})$, where $\mathcal{F} \subseteq \mathcal{T}[t+]$ is finite.*

The following result is proved in the aforementioned paper of Nyikos²⁴.

Theorem 8 – *Let \mathcal{T} be a tree. The following are equivalent.*

- (i) *\mathcal{T} is chain-complete and $\text{MIN}(\mathcal{T})$ is finite.*
- (ii) *The coarse wedge topology of \mathcal{T} is compact and Hausdorff.*

We conclude the current subsection on trees with the following proposition.

²⁴Nyikos, 1997, “Various topologies on trees”, Corollary 3.5.

2. The Szlenk index, trees and operators on Banach spaces over trees

Proposition 7 – Let (T, \leq) be a tree with $ht(T) \leq \omega$.

- (i) Let $S \subseteq T$ be a downwards closed subset of T . Then S is closed in the coarse-wedge topology of T .
- (ii) Let (S, \leq') be a tree and suppose $\phi : S \rightarrow T$ is an order-isomorphism of S onto a downwards closed subset of T . Then ϕ is coarse wedge continuous.

Proof. We first prove (i). Suppose $t \in T \setminus S$. Then $T[t \leq]$ is open in the coarse wedge topology of T since $ht_T(t) < \omega$. Moreover $S \cap T[t \leq] = \emptyset$ since $t \notin S$ and S is downwards closed.

To prove (ii), first note that the sets $T[t \leq]$ and $T \setminus T[t \leq]$, where t varies over all of T , form a subbasis of clopen sets for the coarse wedge topology of T . To establish the continuity of ϕ it therefore suffices to show that $\phi^{-1}(T[t \leq])$ is clopen in S for every $t \in T$. To this end suppose $t \in T$. If $t \notin \phi(S)$ then $\phi^{-1}(T[t \leq]) = \emptyset$ since $\phi(S)$ is downwards closed in T . On the other hand if $t \in \phi(S)$, say $t = \phi(s)$, then $\phi^{-1}(T[t \leq]) = S[s \leq']$ since ϕ is an order isomorphism. \square

2.3 Operators on Banach spaces over trees

Let (T, \leq) be a tree. Define $\Sigma_T : \ell_1(T) \rightarrow \ell_\infty(T)$ by setting

$$\Sigma_T w = \left(\sum_{s \leq t} w(s) \right)_{t \in T}, \quad w \in \ell_1(T).$$

Equivalently, Σ_T is the unique element of $\mathcal{L}(\ell_1(T), \ell_\infty(T))$ satisfying $\Sigma_T e_t = \chi_{T[t \leq]}$ for each $t \in T$, with $\|\Sigma_T\| = 1$ for nonempty T .

Notice that we can state some existing universality results in terms of operators of the form Σ_T . For instance, taking T to be the set of natural numbers \mathbb{N} equipped with its usual order \leq , the operator Σ_T is the aforementioned universal non-weakly compact operator of Lindenstrauss and Pełczyński (Theorem 3 on p. 167). Moreover, taking T to instead be the set of natural numbers \mathbb{N} equipped with the trivial order $=$ yields Σ_T as the formal identity operator from ℓ_1 to ℓ_∞ , shown by Johnson to be universal for the class of non-compact operators (Theorem 4 on p. 167). Amongst the outcomes of the current paper is that we add to the collection of trees (T, \leq) for which the corresponding operator Σ_T is universal for the complement of some operator ideal.

We shall use the following proposition to determine whether Σ_T factors through T , for certain trees (T, \leq) and operators T .

Proposition 8 – Let (T, \leq) be a tree, X and Y Banach spaces and $T \in \mathcal{L}(X, Y)$. The following are equivalent:

- (i) Σ_T factors through T .

(ii) There exist $\delta > 0$ and families $(x_t)_{t \in \mathcal{T}} \subseteq B_X$ and $(x_t^*)_{t \in \mathcal{T}} \subseteq T^*B_{Y^*}$ such that

$$\langle x_t^*, x_s \rangle = \begin{cases} \langle x_s^*, x_s \rangle \geq \delta & \text{if } s \leq t \\ 0 & \text{if } s \not\leq t \end{cases}, \quad s, t \in \mathcal{T}. \quad (7)$$

Proof. First suppose that (i) holds and that $U \in \mathcal{L}(\ell_1(\mathcal{T}), X)$ and $V \in \mathcal{L}(Y, \ell_\infty(\mathcal{T}))$ are such that $VTU = \Sigma_{\mathcal{T}}$. For each $t \in \mathcal{T}$ let f_t^* be the element of $\ell_\infty(\mathcal{T})^*$ satisfying $\langle f_t^*, z \rangle = \|V\|^{-1}z(t)$ for every $z \in \ell_\infty(\mathcal{T})$. For each $s, t \in \mathcal{T}$ let $x_s = \|Ue_s\|^{-1}Ue_s \in B_X$ and $x_t^* = T^*V^*f_t^* \in T^*B_{Y^*}$. Then for $s, t \in \mathcal{T}$ we have

$$\begin{aligned} \langle x_t^*, x_s \rangle &= \langle T^*V^*f_t^*, \|Ue_s\|^{-1}Ue_s \rangle \\ &= \|Ue_s\|^{-1} \langle f_t^*, VTUe_s \rangle \\ &= \|Ue_s\|^{-1} \langle f_t^*, \Sigma_{\mathcal{T}}e_s \rangle \\ &= \begin{cases} \|Ue_s\|^{-1} \|V\|^{-1} \geq (\|U\| \|V\|)^{-1} & \text{if } s \leq t \\ 0 & \text{if } s \not\leq t \end{cases} \end{aligned}$$

hence (i) implies (ii).

Now suppose that (ii) holds and let $\delta > 0$, $(x_t)_{t \in \mathcal{T}} \subseteq B_X$ and $(x_t^*)_{t \in \mathcal{T}} \subseteq T^*B_{Y^*}$ be such that Equation (7) holds. Define $U: \ell_1(\mathcal{T}) \rightarrow X$ by setting

$$Uw = \sum_{t \in \mathcal{T}} \frac{w(t)}{\langle x_t^*, x_t \rangle} x_t$$

for each $w \in \ell_1(\mathcal{T})$. Then U is well-defined, linear and continuous with $\|U\| \leq \delta^{-1}$. For each $t \in \mathcal{T}$ choose $v_t^* \in B_{Y^*}$ such that $T^*v_t^* = x_t^*$. The map $V: Y \rightarrow \ell_\infty(\mathcal{T})$ given by setting $Vy = (\langle v_t^*, y \rangle)_{t \in \mathcal{T}}$ for each $y \in Y$ is well-defined, linear and continuous with $\|V\| \leq 1$. To complete the proof we show that $VTU = \Sigma_{\mathcal{T}}$. To this end note that for $s \in \mathcal{T}$ we have

$$\begin{aligned} VTUe_s &= VT(\langle x_s^*, x_s \rangle^{-1}x_s) = (\langle x_s^*, x_s \rangle^{-1} \langle v_t^*, Tx_s \rangle)_{t \in \mathcal{T}} \\ &= (\langle x_s^*, x_s \rangle^{-1} \langle x_t^*, x_s \rangle)_{t \in \mathcal{T}} = \chi_{\mathcal{T}[s \leq \cdot]}, \end{aligned}$$

hence $VTU = \Sigma_{\mathcal{T}}$. □

The following result may be proved by an appeal to Proposition 8, but an equally easy direct proof is possible (we omit the details).

Proposition 9 – Let (\mathcal{S}, \leq') and (\mathcal{T}, \leq) be trees and suppose that \mathcal{S} is order isomorphic to a subtree of \mathcal{T} . Then $\Sigma_{\mathcal{S}}$ factors through $\Sigma_{\mathcal{T}}$.

Notice that if \mathcal{I} and \mathcal{J} are operator ideals and $T \in \mathcal{I} \cap \mathcal{C}\mathcal{I}$ is universal for $\mathcal{C}\mathcal{I}$, then every universal element of $\mathcal{C}\mathcal{I}$ belongs to \mathcal{J} . In particular, it is a consequence

2. The Szlenk index, trees and operators on Banach spaces over trees

of the following proposition that if \mathcal{S} is an operator ideal and \mathcal{T} is a tree such that $\Sigma_{\mathcal{T}}$ is universal for $\mathcal{C}\mathcal{S}$, then any operator that is universal for $\mathcal{C}\mathcal{S}$ must be strictly singular.

Proposition 10 – *Let (\mathcal{T}, \leq) be a tree. Then:*

- (i) $\Sigma_{\mathcal{T}}$ is strictly singular.
- (ii) $\Sigma_{\mathcal{T}}$ is weakly compact if and only if \mathcal{T} is well-founded.
- (iii) $\Sigma_{\mathcal{T}}$ is compact if and only if \mathcal{T} is finite, if and only if $\Sigma_{\mathcal{T}}$ is finite rank.

Suppose (\mathcal{T}, \leq) is a tree. For the purposes of proving Proposition 10 we now recall the definition of the *James tree space* of \mathcal{T} , denoted $J(\mathcal{T})$, which is the completion of $c_{00}(\mathcal{T})$ with respect to the norm $\|\cdot\|_{J(\mathcal{T})}$ on $c_{00}(\mathcal{T})$ that is defined by setting

$$\|x\|_{J(\mathcal{T})} = \sup \left\{ \left(\sum_{i=1}^k \left| \sum_{t \in \mathcal{S}_i} x(t) \right|^2 \right)^{1/2} \mid \mathcal{S}_1, \dots, \mathcal{S}_k \subseteq \mathcal{T} \text{ pairwise disjoint intervals} \right\}$$

for each $x \in c_{00}(\mathcal{T})$. The formal identity map from $(c_{00}(\mathcal{T}), \|\cdot\|_{\ell_1(\mathcal{T})})$ to $(c_{00}(\mathcal{T}), \|\cdot\|_{J(\mathcal{T})})$ is norm continuous and therefore admits a (unique) continuous linear extension $A_{\mathcal{T}} \in \mathcal{L}(\ell_1(\mathcal{T}), J_2(\mathcal{T}))$. Moreover, the linear map $x \mapsto (\sum_{s \leq t} x(s))_{t \in \mathcal{T}}$ from $c_{00}(\mathcal{T})$ to $\ell_{\infty}(\mathcal{T})$ is norm continuous with respect to the norm $\|\cdot\|_{J(\mathcal{T})}$ on $c_{00}(\mathcal{T})$, and thus extends (uniquely) to some $B_{\mathcal{T}} \in \mathcal{L}(J(\mathcal{T}), \ell_{\infty}(\mathcal{T}))$. Since $\Sigma_{\mathcal{T}} = B_{\mathcal{T}}A_{\mathcal{T}}$ we have that $\Sigma_{\mathcal{T}}$ factors through the James tree space $J(\mathcal{T})$.

Remark 3 – For an introduction to properties of James tree spaces we refer the interested reader to Brackebusch (1988).

Proof (Proof of Proposition 10). Assertion (i) of the proposition follows from the fact that the domain of $\Sigma_{\mathcal{T}}$, namely $\ell_1(\mathcal{T})$, is ℓ_1 -saturated, whilst $\Sigma_{\mathcal{T}}$ has already been seen above to factor through the ℓ_2 -saturated space $J(\mathcal{T})$. (See Lemma 2 and the final remark of Hagler and Odell (1978) for details of the proof that $J(\mathcal{T})$ is ℓ_2 -saturated.)

For (ii), the assertion that $\Sigma_{\mathcal{T}}$ is weakly compact whenever \mathcal{T} is well-founded follows from the aforementioned fact that $\Sigma_{\mathcal{T}}$ factors through the James tree space $J(\mathcal{T})$ of \mathcal{T} and the fact that $J(\mathcal{T})$ is reflexive if and only if \mathcal{T} is well-founded. The proof of this latter fact is obtained via a straightforward transfinite induction on $\rho(\mathcal{T})$, using the following facts: an ℓ_2 -direct sum of a family of reflexive spaces is reflexive; and, for a rooted tree \mathcal{T} , the Banach space $\left(\bigoplus_{t \in \text{MIN}(\mathcal{T}^*)} J(\mathcal{T}[t \leq]) \right)_{\ell_2}$ is isometrically isomorphic to a codimension 1 subspace of $J(\mathcal{T})$; the remaining details are omitted. On the other hand, if \mathcal{T} is not well-founded then \mathcal{T} contains a path order-isomorphic to \mathbb{N} equipped with its usual order \leq . It follows then by Proposition 9 that $\Sigma_{\mathcal{T}}$ factors the universal non-weakly compact operator of Lindenstrauss and Pełczyński (Theorem 3), hence $\Sigma_{\mathcal{T}}$ fails to be weakly compact whenever \mathcal{T} is not well founded.

To prove (iii), first note that if \mathcal{T} is finite then the codomain is the finite-dimensional space $\ell_\infty(\mathcal{T})$, hence $\Sigma_{\mathcal{T}}$ is finite rank and therefore compact. Conversely, if \mathcal{T} is infinite then the set $\{\Sigma_{\mathcal{T}}e_t \mid t \in \mathcal{T}\}$ is an infinite 1-separated subset of $\Sigma_{\mathcal{T}}B_{\ell_1(\mathcal{T})}$, hence in this case $\Sigma_{\mathcal{T}}$ is non-compact, hence non-finite rank. \square

We now establish a connection between the rank $\rho(\mathcal{T})$ and the Szlenk indices of $\Sigma_{\mathcal{T}}$ in the particular case that \mathcal{T} is blossomed. The proof of the following proposition is similar to the last part of the proof Proposition 6.2 of Beanland and Causey (2018).

Proposition 11 – *Let (\mathcal{T}, \leq) be a blossomed tree and $\epsilon \in (0, 1)$. Then $Sz(\Sigma_{\mathcal{T}^*}, \epsilon) \geq \rho(\mathcal{T})$.*

Proof. Fix $\epsilon \in (0, 1)$, let t_\emptyset denote the root of \mathcal{T} and let $B = \Sigma_{\mathcal{T}^*}^* B_{\ell_\infty(\mathcal{T}^*)}$. For each $t \in \mathcal{T}^*$ let $f_t^* \in \ell_\infty(\mathcal{T}^*)^*$ be the evaluation functional of $\ell_\infty(\mathcal{T}^*)$ at t and let $g_t^* := \Sigma_{\mathcal{T}^*}^* f_t^* \in B$. Let $g_{t_\emptyset}^* := 0 \in B$. Note that $\{g_t^* \mid t \in \mathcal{T}\}$ is ϵ -separated, for if $s, t \in \mathcal{T}$ are such that $s \notin \mathcal{T}[t \leq]$ then $\|g_t^* - g_s^*\| \geq \langle g_t^* - g_s^*, e_t \rangle = 1 > \epsilon$. For each $t \in \mathcal{T} \setminus \text{MAX}(\mathcal{T})$ let $(t_m)_{m=0}^\infty$ be an enumeration of $\mathcal{T}[t+]$ with $(\rho_{\mathcal{T}}(t_m))_{m=0}^\infty$ non-decreasing. For every $x \in \ell_1(\mathcal{T}^*)$ and $t \in \mathcal{T} \setminus \text{MAX}(\mathcal{T})$ we have $\langle g_{t_m}^*, x \rangle = \sum_{s \leq t_m} x(s) \rightarrow \sum_{s \leq t} x(s) = \langle g_t^*, x \rangle$ as $m \rightarrow \infty$, so that $(g_{t_m}^*)_{m=0}^\infty$ is weak*-convergent to g_t^* .

We will show by transfinite induction that

$$\{g_t^* \mid t \in \mathcal{T}^{(\xi)} \setminus \mathcal{T}^{(\xi+1)}\} \subseteq s_\epsilon^\xi(B). \quad (8)$$

for every $\xi < \rho(\mathcal{T})$. Equation (8) is trivially true for $\xi = 0$. Suppose $\zeta \in (0, \rho(\mathcal{T}))$ is such that Equation (8) holds for every $\xi < \zeta$; to complete the induction we show that $\{g_t^* \mid t \in \mathcal{T}^{(\zeta)} \setminus \mathcal{T}^{(\zeta+1)}\} \subseteq s_\epsilon^\zeta(B)$. To this end suppose $t \in \mathcal{T}^{(\zeta)} \setminus \mathcal{T}^{(\zeta+1)}$. By the inductive hypothesis and the fact that $(\rho_{\mathcal{T}}(t_m))_{m=1}^\infty$ is non-decreasing we have that $(g_{t_l}^*)_{l=m}^\infty \subseteq s_\epsilon^{\rho_{\mathcal{T}}(t_m)}(B)$. Thus $g_t^* \in s_\epsilon^{\rho_{\mathcal{T}}(t_m)}(B)$ as the latter set is weak*-closed. This yields $g_t^* \in s_\epsilon^{\rho_{\mathcal{T}}(t_m)+1}(B)$ since $\|g_t^* - g_{t_l}^*\| > \epsilon$ for all $l \in [m, \omega)$. It follows that $g_t^* \in \bigcap_{m < \omega} s_\epsilon^{\rho_{\mathcal{T}}(t_m)+1}(B) = s_\epsilon^{\rho_{\mathcal{T}}(t)}(B) = s_\epsilon^\zeta(B)$, as required. With the induction now complete, taking $\xi = \rho(\mathcal{T}) - 1$ in Equation (8) yields $g_{t_\emptyset}^* \in s_\epsilon^{\rho(\mathcal{T})-1}(B)$, from which the proposition follows. \square

3 Absolutely convex sets of large Szlenk index

This section is devoted to proving our key technical result, Theorem 9, from which the main results of the paper (stated in the Introduction) and a number of other results in subsequent sections of the paper are derived. In order to state Theorem 9 we introduce the following notation. For a family $\mathfrak{T} = ((\mathcal{T}_n, \leq_n))_{n < \omega}$, where each (\mathcal{T}_n, \leq_n) is a rooted tree, define $\llbracket \mathfrak{T} \rrbracket := \{\emptyset\} \cup \bigcup_{n < \omega} (\{n\} \times \mathcal{T}_n^*)$, so that $\llbracket \mathfrak{T} \rrbracket$ is a rooted tree when equipped with the order $\leq_{\mathfrak{T}}$ on $\llbracket \mathfrak{T} \rrbracket$ defined by setting $\emptyset \leq_{\mathfrak{T}} t$ for all $t \in \llbracket \mathfrak{T} \rrbracket$ and $(n_1, t_1) \leq_{\mathfrak{T}} (n_2, t_2)$ if and only if $n_1 = n_2$ and $t_1 \leq_{n_1} t_2$.

3. Absolutely convex sets of large Szlenk index

Theorem 9 – Let X be a Banach space, $K \subseteq X^*$ a non-empty, absolutely convex, weak*-compact set, $\delta, \theta > 0$ positive real numbers, $(\epsilon_n)_{n < \omega}$ a family of positive real numbers, $(\xi_n)_{n < \omega}$ a family of countable ordinals such that $s_{\epsilon_n}^{\xi_n}(K) \neq \emptyset$ for all $n < \omega$, and $\mathfrak{T} = ((T_n, \leq_n))_{n < \omega}$ a family of countable, well-founded, rooted trees such that $\rho(T_n) \leq \xi_n + 1$ for all $n < \omega$.

(i) There exist families $(x_t^*)_{t \in \llbracket \mathfrak{T} \rrbracket^*} \subseteq K$ and $(x_t)_{t \in \llbracket \mathfrak{T} \rrbracket^*} \subseteq S_X$ such that

$$\langle x_t^*, x_s \rangle = \begin{cases} \langle x_s^*, x_s \rangle > \frac{\epsilon_n}{8+\theta} & \text{if } s \leq_{\mathfrak{T}} t \in \{n\} \times T_n^* \\ 0 & \text{if } s \not\leq_{\mathfrak{T}} t \end{cases}, \quad s, t \in \llbracket \mathfrak{T} \rrbracket^*, n < \omega. \quad (9)$$

Moreover, for any bijection $\tau : \omega \rightarrow \llbracket \mathfrak{T} \rrbracket^*$ such that $\tau(l) \leq_{\mathfrak{T}} \tau(m)$ implies $l \leq m$ we may choose $(x_t)_{t \in \llbracket \mathfrak{T} \rrbracket^*}$ so that $(x_{\tau(m)})_{m=1}^{\infty}$ is a basic sequence with basis constant not exceeding $1 + \delta$.

(ii) If X^* is norm separable then the families $(x_t^*)_{t \in \llbracket \mathfrak{T} \rrbracket^*}$ and $(x_t)_{t \in \llbracket \mathfrak{T} \rrbracket^*}$ in (i) may be chosen so that $(x_{\tau(m)})_{m=1}^{\infty}$ is shrinking and $(Qx_{\tau(m)})_{m=1}^{\infty}$ is a shrinking basis for X/Z with basis constant not exceeding $1 + \delta$. Here $Z = \bigcap_{t \in \llbracket \mathfrak{T} \rrbracket^*} \ker(x_t^*)$ and $Q : X \rightarrow X/Z$ is the quotient map.

The proof of Theorem 9 shall invoke the following lemma due to G. Lancien²⁵, who established the result for the special case $K = B_{X^*}$ and ζ of the form ω^α for some ordinal α ; the same argument gives the more general statement presented below.

Lemma 1 – Let X be a Banach space, $K \subseteq X^*$ an absolutely convex, weak*-compact set, ζ an ordinal and $\epsilon > 0$. If $s_\epsilon^\zeta(K) \neq \emptyset$ then

$$\forall n < \omega \quad 0 \in s_{\epsilon/2^{n+1}}^{\zeta 2^n}(K). \quad (10)$$

We require the following result, which is Lemma 2.2 of Dilworth et al. (2017).

Lemma 2 – Let X be a Banach space, $\nu > 0$ a real number, F a finite-dimensional subspace of X^* , A a $\frac{\nu}{4+2\nu}$ -net in S_F and $\{y_{f^*} \mid f^* \in A\} \subseteq S_X$ a family such that $\inf\{|f^*(y_{f^*})| \mid f^* \in A\} \geq \frac{4+\nu}{4+2\nu}$. Then for every $x^* \in \{y_{f^*} \mid f^* \in A\}^\perp$ we have $\sup\{|x^*(y)| \mid y \in B_{F^\perp}\} \geq \frac{1}{2+\nu} \|x^*\|$.

We do not know a reference for the following result, so we provide the straightforward proof.

Lemma 3 – Let X be a Banach space and $x^* \in X^*$. Then

$$\{x^* + \epsilon B_{X^*}^\circ + C^\perp \mid \epsilon > 0, C \subseteq X, |C| < \infty\}$$

is a local base for the weak* topology of X^* at x^* .

²⁵Lancien, 1996, “On the Szlenk index and the weak*-dentability index”, p. 67.

Proof. We assume $x^* = 0$, from which the general case follows easily. For $\epsilon > 0$ and finite $C \subseteq X$ we have $\epsilon B_{X^*}^\circ + C^\perp \subseteq \bigcap_{x \in C} \{y^* \in X^* \mid |y^*(x)| < \epsilon\}$, so it suffices to show that $\epsilon B_{X^*}^\circ + C^\perp$ is weak*-open for $\epsilon > 0$ and finite $C \subseteq X$.

Fix $\epsilon > 0$ and a finite set $C \subseteq X$. Set $Y = \text{span}(C)$. Notice that $C^\perp = Y^\perp$. We have

$$\epsilon B_{X^*}^\circ + C^\perp = \{x^* \in X^* \mid \text{dist}(x^*, Y^\perp) < \epsilon\} = Q^{-1}(\epsilon B_{X^*/Y^\perp}^\circ)$$

where $Q : X^* \rightarrow X^*/Y^\perp$ is the canonical quotient map. It is thus enough to show that Q is weak* continuous. This follows from the classic decomposition $Q = S \circ R$, where $R : X^* \rightarrow Y^*$ is the restriction map, which is clearly weak* continuous, and $S : Y^\perp \rightarrow X^*/Y^\perp$, defined by $S(Rx^*) = Qx^*$, is weak* continuous as a bounded linear map between finite dimension normed spaces. \square

Lemma 4 – *Let X be a Banach space, K and L weak*-compact subsets of X^* , ξ an ordinal and $\epsilon > 0$. If $x^* \in s_\epsilon^\xi(K)$ and $y^* \in L$ then $x^* + y^* \in s_\epsilon^\xi(K + L)$.*

Proof. We proceed by transfinite induction on ξ . The assertion of the lemma is true for $\xi = 0$. Suppose that $\zeta > 0$ is an ordinal such that the assertion of the lemma is true for $\xi = \zeta$; we will show that it is true then for $\xi = \zeta + 1$. Let $x^* \in s_\epsilon^{\zeta+1}(K)$ and $y^* \in L$. Since $x^* \in s_\epsilon^\zeta(K)$ it follows from the induction hypothesis that $x^* + y^* \in s_\epsilon^\zeta(K + L)$. Since $x^* \in s_\epsilon^{\zeta+1}(K)$ we have that for any weak*-neighbourhood \mathcal{U} of $x^* + y^*$ there exist $x_1^*, x_2^* \in s_\epsilon^\zeta(K) \cap (-y^* + \mathcal{U})$ such that $\|x_1^* - x_2^*\| > \epsilon$. Since $x_1^* + y^*, x_2^* + y^* \in \mathcal{U} \cap (K + L)$ and $\|(x_1^* + y^*) - (x_2^* + y^*)\| > \epsilon$, we deduce that $x^* + y^* \in s_\epsilon^{\zeta+1}(K + L)$.

Now suppose ζ is a limit ordinal and the assertion of the lemma is true for all $\xi < \zeta$. For $x^* \in s_\epsilon^\zeta(K) = \bigcap_{\xi < \zeta} s_\epsilon^\xi(K)$ and $y^* \in L$ we have

$$x^* + y^* \in \bigcap_{\xi < \zeta} s_\epsilon^\xi(K + L) = s_\epsilon^\zeta(K + L),$$

which completes the induction. \square

The final preliminary result before proving Theorem 9 is the following theorem due to Kadec²⁶ and Klee²⁷; a short proof due to Davis and Johnson (sketched in Davis and Johnson 1973) has been published by Lindenstrauss and Tzafriri²⁸.

Theorem 10 – *Let $(X, \|\cdot\|)$ be a Banach space and $c > 1$ a real number. If X^* is norm separable then X admits a norm $\|\|\cdot\|\|$ such that the following properties hold:*

- (i) *For every $x^* \in X^*$ we have $\|x^*\| \leq \|\|\cdot\|\| \leq c\|x^*\|$.*

²⁶Kadec, 1958, "On strong and weak convergence".

²⁷Klee, 1960, "Mappings into normed linear spaces".

²⁸Lindenstrauss and Tzafriri, 1977, *Classical Banach spaces. I*, p. 13.

3. Absolutely convex sets of large Szlenk index

(ii) If $(x_n^*) \subseteq X^*$ and $x^* \in X^*$ are such that $x_n^* \xrightarrow{w^*} x^*$ and $\|x_n^*\| \rightarrow \|x^*\|$, then $\|x_n^* - x^*\| \rightarrow 0$.

As is implicit in the statement of Theorem 10, when we apply the renorming result Theorem 10 we shall use $\|\cdot\|$ to denote also the corresponding induced norm on duals, subspaces, quotients and operators on X .

Proof (Proof of Theorem 9). We shall first prove (i) without any assumption on the norm density of X^* , then show how to modify the arguments in the proof of (i) to obtain also the assertions of (ii) when X^* is assumed to be norm separable.

Fix $\delta, \theta > 0$ and let $\nu > 0$ be a real number small enough that

$$\frac{1 - 2\nu}{4(2 + \nu)(1 + \nu)} \geq \frac{1}{8 + \theta}. \quad (11)$$

Let $\tau : \omega \rightarrow \llbracket \mathfrak{T} \rrbracket$ be a bijection such that $\tau(l) \leq_{\mathfrak{T}} \tau(m)$ implies $l \leq m$ (c.f. the paragraph preceding Example 1), noting that we necessarily have $\tau(0) = \emptyset$, the root of $\llbracket \mathfrak{T} \rrbracket$. For $0 < m < \omega$ we may write $\tau(m) = (n_m, t_m)$, where $n_m < \omega$ and $t_m \in T_{n_m}^*$. Fix a sequence $(\delta_m)_{m=1}^\infty \subseteq (0, 1)$ of positive real numbers such that $\sum_{m=1}^\infty \delta_m < \infty$ and $\prod_{m=1}^\infty (1 - \delta_m) \geq (1 + \delta)^{-1}$. Proceeding via an induction over $m \in [1, \omega)$, we shall construct families $(f_{\tau(m)}^*)_{m < \omega} \subseteq X^*$ and $(x_{\tau(m)})_{1 \leq m < \omega} \subseteq S_X$ satisfying the following conditions for all $m \in [1, \omega)$:

$$(I) \quad f_{\tau(m)}^* \in s_{\epsilon_{n_m}/2}^{\rho_{T_{n_m}}(t_m)} \left((1 + \nu \sum_{j=1}^m 2^{-j})K \right).$$

(II) For all $i, j \in [1, m]$,

$$\langle f_{\tau(j)}^*, x_{\tau(i)} \rangle = \begin{cases} \langle f_{\tau(i)}^*, x_{\tau(i)} \rangle > \frac{(1-2\nu)\epsilon_{n_i}}{4(2+\nu)} & \text{if } \tau(i) \leq_{\mathfrak{T}} \tau(j) \\ 0 & \text{if } \tau(i) \not\leq_{\mathfrak{T}} \tau(j) \end{cases}. \quad (12)$$

(III) For all $x \in \text{span}\{x_{\tau(l)} \mid 1 \leq l < m\}$ and scalars a we have $\|x + a x_{\tau(m)}\| \geq (1 - \delta_m)\|x\|$.

Since for $0 < m < \omega$ we have $s_{\epsilon_{n_m}/2}^{\rho_{T_{n_m}}(t_m)} \left((1 + \nu \sum_{j=1}^m 2^{-j})K \right) \subseteq (1 + \nu)K$, once the induction is complete the first assertion of (i) then follows from Equation (11), (I) and (II) by taking $x_t^* = \frac{1}{1+\nu} f_t^*$ for each $t \in \llbracket \mathfrak{T} \rrbracket^*$.

The second assertion of (i) follows from the Grunblum criterion²⁹ and the fact that, by (III), for $1 \leq l < m < \omega$ and scalars a_1, \dots, a_m we have

$$\left\| \sum_{q=1}^l a_q x_{\tau(q)} \right\| \leq \frac{1}{\prod_{q=l+1}^m (1 - \delta_q)} \left\| \sum_{q=1}^m a_q x_{\tau(q)} \right\| \leq (1 + \delta) \left\| \sum_{q=1}^m a_q x_{\tau(q)} \right\|.$$

For each $n < \omega$ let o_n denote the root of \mathcal{T}_n . It follows from Equation (10) that

$$0 \in \bigcap_{n < \omega} s_{\epsilon_n/2}^{\xi_n}(K) \subseteq \bigcap_{n < \omega} s_{\epsilon_n/2}^{\rho(\mathcal{T}_n)-1}(K) = \bigcap_{n < \omega} s_{\epsilon_n/2}^{\rho_{\mathcal{T}_n}(o_n)}(K). \quad (13)$$

Define $f_{\tau(0)}^* := 0 \in \bigcap_{n < \omega} s_{\epsilon_n/2}^{\xi_n}(K)$. Since

$$f_{\tau(0)}^* \in s_{\epsilon_{n_1}/2}^{\rho_{\mathcal{T}_{n_1}}(o_{n_1})}(K) \subseteq s_{\epsilon_{n_1}/2}^{\rho_{\mathcal{T}_{n_1}}(t_1)+1}(K) = s_{\epsilon_{n_1}/2} \left(s_{\epsilon_{n_1}/2}^{\rho_{\mathcal{T}_{n_1}}(t_1)}(K) \right),$$

it follows from the definition of the derivation $s_{\epsilon_{n_1}/2}$ that there is $f_{\tau(1)}^* \in s_{\epsilon_{n_1}/2}^{\rho_{\mathcal{T}_{n_1}}(t_1)}(K)$ such that $\|f_{\tau(1)}^*\| = \|f_{\tau(1)}^* - f_{\tau(0)}^*\| > \epsilon_{n_1}/4$. Choose $x_{\tau(1)} \in S_X$ so that

$$\langle f_{\tau(1)}^*, x_{\tau(1)} \rangle > \frac{\epsilon_{n_1}}{4}.$$

It is readily checked that (I)–(III) hold for $m = 1$.

Fix $k \in [1, \omega)$ and suppose that the points $f_{\tau(m)}^* \in s_{\epsilon_{n_m}/2}^{\rho_{\mathcal{T}_{n_m}}(t_m)} \left((1 + \nu \sum_{j=1}^m 2^{-j})K \right)$ and $x_{\tau(m)} \in S_X$ have been defined for $1 \leq m \leq k$ in such a way that properties (I)–(III) are satisfied for $1 \leq m \leq k$. Let $(k+1)^-$ denote the unique ordinal less than $k+1$ and such that $\tau((k+1)^-) = \tau(k+1)^-$. To carry out the inductive step of the proof we show how to construct $f_{\tau(k+1)}^* \in s_{\epsilon_{n_{k+1}}/2}^{\rho_{\mathcal{T}_{n_{k+1}}}(t_{k+1})} \left((1 + \nu \sum_{j=1}^{k+1} 2^{-j})K \right)$ and $x_{\tau(k+1)} \in S_X$ so that (I)–(III) are satisfied for $1 \leq m \leq k+1$. Our first task will be to define $f_{\tau(k+1)}^*$ as a point inside a certain weak*-neighbourhood of $f_{\tau(k+1)^-}^*$ and then show that (I) holds for $m = k+1$. To this end let G be a finite δ_{k+1} -net in $S_{\text{span}\{x_{\tau(i)} \mid 1 \leq i \leq k\}}$ and for each $g \in G$ let $h_g^* \in X^*$ be such that $\langle h_g^*, g \rangle = 1$. Set $F = \text{span} \left(\{f_{\tau(i)}^* \mid 1 \leq i \leq k\} \cup \{h_g^* \mid g \in G\} \right) \subseteq X^*$, let A be a finite $\frac{\nu}{4+2\nu}$ -net in S_F and let $\{y_{f^*} \mid f^* \in A\} \subseteq S_X$ be such that $f^*(y_{f^*}) \geq \frac{4+\nu}{4+2\nu}$ for each $f^* \in A$. Let

$$\mathcal{U}_1 = \bigcap_{i=1}^k \left\{ x^* \in X^* \mid |\langle x^* - f_{\tau(k+1)^-}^*, x_{\tau(i)} \rangle| < \frac{2^{-k-4}\nu(1-2\nu)\epsilon_{n_i}}{k(2+\nu)(1+\nu)} \right\} \quad \text{and}$$

$$\mathcal{U}_2 = f_{\tau(k+1)^-}^* + \frac{\nu\epsilon_{n_{k+1}}}{4(2+\nu)} B_{X^*}^\circ + \left(\{x_{\tau(i)} \mid 1 \leq i \leq k\} \cup \{y_{f^*} \mid f^* \in A\} \right)^\perp.$$

Note that \mathcal{U}_2 is a weak*-open neighbourhood of $f_{\tau(k+1)^-}^*$ by Lemma 3, hence $\mathcal{U} := \mathcal{U}_1 \cap \mathcal{U}_2$ is a weak*-open neighbourhood of $f_{\tau(k+1)^-}^*$. On the one hand, if $(k+1)^- = 0$ then $f_{\tau(k+1)^-}^* = 0$, hence by Equation (13) in this case we have

$$f_{\tau(k+1)^-}^* = 0 \in s_{\epsilon_{n_{k+1}}/2}^{\rho_{\mathcal{T}_{n_{k+1}}}(o_{n_{k+1}})}(K) \subseteq s_{\epsilon_{n_{k+1}}/2}^{\rho_{\mathcal{T}_{n_{k+1}}}(o_{n_{k+1}})} \left((1 + \nu \sum_{j=1}^k 2^{-j})K \right)$$

(Cont. next page)

3. Absolutely convex sets of large Szlenk index

$$\subseteq s_{\epsilon_{n_{k+1}}/2}^{\rho_{\mathcal{T}_{n_{k+1}}}(t_{k+1})+1} \left((1 + \nu \sum_{j=1}^k 2^{-j})K \right). \quad (14)$$

On the other hand, if $(k+1)^- \neq 0$ then $n_{(k+1)^-} = n_{k+1}$ and $t_{(k+1)^-} = t_{k+1}^- \in \mathcal{T}_{n_{k+1}}$. So by the hypothesis that (I) holds for $1 \leq m \leq k$, we have

$$\begin{aligned} f_{\tau(k+1)^-}^* &\in s_{\epsilon_{n_{k+1}}/2}^{\rho_{\mathcal{T}_{n_{k+1}}}(t_{k+1}^-)} \left((1 + \nu \sum_{j=1}^{(k+1)^-} 2^{-j})K \right) \subseteq s_{\epsilon_{n_{k+1}}/2}^{\rho_{\mathcal{T}_{n_{k+1}}}(t_{k+1}^-)} \left((1 + \nu \sum_{j=1}^k 2^{-j})K \right) \\ &\subseteq s_{\epsilon_{n_{k+1}}/2}^{\rho_{\mathcal{T}_{n_{k+1}}}(t_{k+1})+1} \left((1 + \nu \sum_{j=1}^k 2^{-j})K \right). \end{aligned} \quad (15)$$

It follows from Equations (14) and (15), and the definition of the derivation s_e^ξ for $\epsilon > 0$ and $\xi \in \mathbf{Ord}$ that there exists $u^* \in \mathcal{U} \cap s_{\epsilon_{n_{k+1}}/2}^{\rho_{\mathcal{T}_{n_{k+1}}}(t_{k+1})} \left((1 + \nu \sum_{j=1}^k 2^{-j})K \right)$ such that $\|f_{\tau(k+1)^-}^* - u^*\| > \epsilon_{n_{k+1}}/4$. Define

$$f_{\tau(k+1)}^* := u^* - \sum_{l=1}^k \frac{\langle u^* - f_{\tau(k+1)^-}^*, x_{\tau(l)} \rangle}{\langle f_{\tau(l)}^*, x_{\tau(l)} \rangle} (f_{\tau(l)}^* - f_{\tau(l)^-}^*).$$

Since $u^* \in \mathcal{U}_1$ and since $f_{\tau(l)}^* - f_{\tau(l)^-}^* \in 2(1 + \nu)K$ for $1 \leq l \leq k$ (by the assumption that (I) holds for $1 \leq m \leq k$), it follows from the definition of $f_{\tau(k+1)}^*$ that we have $f_{\tau(k+1)}^* - u^* \in cK$, where $c > 0$ is a scalar that may be taken to satisfy

$$\begin{aligned} c &\leq \sum_{l=1}^k \frac{|\langle u_{\tau(k+1)}^* - f_{\tau(k+1)^-}^*, x_{\tau(l)} \rangle|}{|\langle f_{\tau(l)}^*, x_{\tau(l)} \rangle|} 2(1 + \nu) \\ &\leq \sum_{l=1}^k \frac{2^{-k-4} \nu (1 - 2\nu) \epsilon_{n_l}}{k(2 + \nu)(1 + \nu)} \frac{4(2 + \nu)}{(1 - 2\nu) \epsilon_{n_l}} 2(1 + \nu) \\ &= \sum_{l=1}^k \frac{2^{-k-1} \nu}{k} \\ &= \nu 2^{-k-1}. \end{aligned} \quad (16)$$

An appeal to Lemma 4 yields

$$f_{\tau(k+1)}^* = u^* + (f_{\tau(k+1)}^* - u^*) \in s_{\epsilon_{n_{k+1}}/2}^{\rho_{\mathcal{T}_{n_{k+1}}}(t_{k+1})} \left(\left(1 + \nu \sum_{j=1}^{k+1} 2^{-j} \right) K \right)$$

hence (I) holds for $m = k + 1$.

We now show how to define $x_{\tau(k+1)}$ and then verify that (II) and (III) hold for $m = k + 1$. Since $u^* \in \mathcal{U}_2$ we may write $u^* = f_{\tau(k+1)-}^* + y^* + x^*$, where $\|y^*\| < \frac{\nu\epsilon_{n_{k+1}}}{4(2+\nu)}$ and $x^* \in (\{x_{\tau(i)} \mid 1 \leq i \leq k\} \cup \{y_{f^*} \mid f^* \in A\})^\perp$. Since

$$\|x^*\| \geq \|u^* - f_{\tau(k+1)-}^*\| - \|y^*\| > \frac{\epsilon_{n_{k+1}}}{4} - \frac{\nu\epsilon_{n_{k+1}}}{4(2+\nu)} > \frac{(1-\nu)\epsilon_{n_{k+1}}}{4},$$

an application of Lemma 2 with F, A and $\{y_{f^*} \mid f^* \in A\} \subseteq S_X$ as defined above in the current proof yields $y \in S_{F_\perp}$ such that

$$\langle x^*, y \rangle > \frac{1}{2+\nu} \frac{(1-\nu)\epsilon_{n_{k+1}}}{4}.$$

Set $x_{\tau(k+1)} = |\langle u^*, y \rangle| \langle u^*, y \rangle^{-1} y$, so that $x_{\tau(k+1)} \in S_{F_\perp} \subseteq S_X$ and

$$\begin{aligned} \langle u^*, x_{\tau(k+1)} \rangle &= |\langle u^*, y \rangle| = |\langle u^* - f_{\tau(k+1)-}^*, y \rangle| = |\langle x^* + y^*, y \rangle| \\ &> \frac{(1-\nu)\epsilon_{n_{k+1}}}{4(2+\nu)} - \frac{\nu\epsilon_{n_{k+1}}}{4(2+\nu)} = \frac{(1-2\nu)\epsilon_{n_{k+1}}}{4(2+\nu)}. \end{aligned}$$

We now show that (II) holds for $m = k + 1$. By the induction hypothesis, we need to prove the case where at least one of i and j is equal to $k + 1$. Since $x_{\tau(k+1)} \in S_{F_\perp} \subseteq \bigcap_{j=1}^k \ker(f_{\tau(j)}^*)$ we have $\langle f_{\tau(j)}^*, x_{\tau(k+1)} \rangle = 0$ for $1 \leq j \leq k$. Moreover it is clear from the definition of $f_{\tau(k+1)}^*$ and the fact that $x_{\tau(k+1)} \in \bigcap_{j=1}^k \ker(f_{\tau(j)}^*)$ that

$$\langle f_{\tau(k+1)}^*, x_{\tau(k+1)} \rangle = \langle u^*, x_{\tau(k+1)} \rangle - 0 = \langle u^*, x_{\tau(k+1)} \rangle > \frac{(1-2\nu)\epsilon_{n_{k+1}}}{4(2+\nu)}.$$

Since (II) holds for $1 \leq m \leq k$, if $l, i \in [1, k]$ then

$$\langle f_{\tau(l)}^* - f_{\tau(l)-}^*, x_{\tau(i)} \rangle = \begin{cases} \langle f_{\tau(i)}^*, x_{\tau(i)} \rangle, & \text{if } i = l \\ 0, & \text{if } i \neq l \end{cases} \quad (17)$$

It follows that if $1 \leq i \leq k$ then

$$\begin{aligned} \langle f_{\tau(k+1)}^*, x_{\tau(i)} \rangle &= \langle u^*, x_{\tau(i)} \rangle - \frac{\langle u^* - f_{\tau(k+1)-}^*, x_{\tau(i)} \rangle}{\langle f_{\tau(i)}^*, x_{\tau(i)} \rangle} \langle f_{\tau(i)}^*, x_{\tau(i)} \rangle \\ &= \langle f_{\tau(k+1)-}^*, x_{\tau(i)} \rangle \\ &= \begin{cases} \langle f_{\tau(i)}^*, x_{\tau(i)} \rangle > \frac{(1-2\nu)\epsilon_{n_i}}{4(2+\nu)} & \text{if } \tau(i) \leq_{\mathfrak{T}} \tau(k+1) \\ 0 & \text{if } \tau(i) \not\leq_{\mathfrak{T}} \tau(k+1) \end{cases}, \end{aligned}$$

3. Absolutely convex sets of large Szlenk index

hence (II) holds for $m = k + 1$.

Finally, we show that (III) holds for $m = k + 1$. Let a be a scalar and, to avoid triviality, let $x \in \text{span}\{x_{\tau(i)} \mid 1 \leq i \leq k\}$ be nonzero. Let $g_x \in G$ be such that $\| \|x\|^{-1}x - g_x \| \leq \delta_{k+1}$. Since $\langle h_{g_x}^*, g_x \rangle = 1$ and $x_{\tau(k+1)} \in \ker(h_{g_x}^*)$ we have

$$\begin{aligned} \|x + ax_{\tau(k+1)}\| &\geq |\langle h_{g_x}^*, x + ax_{\tau(k+1)} \rangle| \\ &= |\langle h_{g_x}^*, \|x\|^{-1}x \rangle| \|x\| \\ &\geq \left(|\langle h_{g_x}^*, g_x \rangle| - |\langle h_{g_x}^*, \|x\|^{-1}x - g_x \rangle| \right) \|x\| \\ &\geq (1 - \delta_{k+1}) \|x\|, \end{aligned}$$

which completes the proof of part (i) of the theorem.

We now prove (ii). To this end suppose X^* is norm separable and let $(z_m^*)_{m=1}^\infty$ a norm dense sequence in X^* . To see that $(x_{\tau(m)})_{m=1}^\infty$ may be chosen to be a shrinking basis, we modify the proof of (i) by extending the list of conditions (I)–(III) to include the following fourth condition:

$$(IV) \quad x_{\tau(m)} \in \bigcap_{j=1}^{m-1} \ker(z_j^*).$$

In the inductive construction involving the verification of properties (I)–(III), we amend the argument to ensure that (IV) holds for all $m \in [1, \omega)$ as follows. For the base step we require no change in the argument since $\bigcap_{j=1}^0 \ker(z_j^*) = X$. For the inductive step we change the definition of F so that

$$F = \text{span}\left(\{f_{\tau(i)}^* \mid 1 \leq i \leq k\} \cup \{h_g^* \mid g \in G\} \cup \{z_j^* \mid 1 \leq j \leq k\}\right).$$

Since $x_{\tau(k+1)}$ is defined so that $x_{\tau(k+1)} \in S_{F^\perp}$, the induction yields that (IV) holds for all $m \in [1, \omega)$. With this modification it is now easy to see that $(x_{\tau(m)})_{m=1}^\infty$ is shrinking. Indeed, let $f^* \in [(x_{\tau(m)})_{m=1}^\infty]^*$ and fix $\epsilon > 0$. Let $\tilde{f}^* \in X^*$ be an extension of f^* to X and let $N < \omega$ be such that $\|\tilde{f}^* - z_N^*\| < \epsilon$. By (IV) we have $\langle z_N^*, x_{\tau(m)} \rangle = 0$ for all $m > N$, hence $m > N$ implies

$$\|f^*|_{[x_{\tau(m)}]_{m>N}}\| = \|\tilde{f}^*|_{[x_{\tau(m)}]_{m>N}}\| \leq \|(\tilde{f}^* - z_N^*)|_{[x_{\tau(m)}]_{m>N}}\| + \|z_N^*|_{[x_{\tau(m)}]_{m>N}}\| < \epsilon.$$

In particular, $\lim_{M \rightarrow \infty} \|f^*|_{[x_{\tau(m)}]_{m>M}}\| = 0$. As $f^* \in [(x_{\tau(m)})_{m=1}^\infty]^*$ was chosen arbitrarily, $(x_{\tau(m)})_{m=1}^\infty$ is shrinking by Proposition 3.2.8 of Albiac and Kalton (2016).

We now show how to modify the proof of (i) further so that $(Qx_{\tau(m)})_{m=1}^\infty$ is a shrinking basis for X/Z with basis constant not exceeding $1 + \delta$. In a similar spirit to the proof of Proposition 3.5 of Lancien (1996), the main idea is to modify the proof of (i) to incorporate the arguments from Johnson and Rosenthal's proof of Theorem III.1 of Johnson and Rosenthal (1972). To this end let $\|\cdot\|$ be an equivalent norm on X such that properties (i) and (ii) of Theorem 10 hold with $c = (1 + \delta)^{1/2}$ and let $(z_p)_{p=1}^\infty$ be a norm dense sequence in $S_{(X, \|\cdot\|)}$. Fix a sequence

$(\delta'_m)_{m=1}^\infty \subseteq (0, 1) \subseteq \mathbb{R}$ such that $\sum_{m=1}^\infty \delta'_m < \infty$ and $\prod_{m=1}^\infty (1 - \delta'_m) \geq (1 + \delta)^{-1/2}$. At the k th stage of the inductive construction, after having defined the new $f_{\tau(k)}^*$, we set $v_k := \|\|f_{\tau(k)}^* - f_{\tau(k)}^*\|^{-1} (f_{\tau(k)}^* - f_{\tau(k)}^*)$. We modify the inductive construction in proof of (i) to include the construction of a strictly increasing sequence $(p_m)_{m=0}^\infty \subseteq \omega$ such that the following additional conditions hold for $m \in [1, \omega)$:

(V) For each $v^* \in [(v_j)_{j=1}^m]^*$ with $\|\|v^*\| = 1$ there is a natural number $p \leq p_m$ such that $|\langle v, z_p \rangle - \langle v^*, v \rangle| \leq \|\|v\| \delta'_m / 3$ for all $v \in [(v_j)_{j=1}^m]$.

(VI) $|\langle v_m, z_p \rangle| < \delta'_m / 3$ for all z_p with $p \leq p_{m-1}$.

At the base step of the induction we set $p_0 = 0$ and use Helly's theorem (or Goldstine's theorem), the density of $(z_p)_{p=1}^\infty$ in $S_{(X, \|\|\cdot\|)}$ and the total boundedness of $S_{(\{(v_1\}, \|\|\cdot\|)}$ and $S_{(\{(v_1\}, \|\|\cdot\|)}$ to obtain also $p_1 > p_0$ large enough that (V) holds for $m = 1$ (we leave the straightforward details to the reader). Since p_0 is defined to be 0, which is not in the index set of the sequence $(z_p)_{p=1}^\infty$, (VI) is true for $m = 1$.

At the inductive step of the modified construction we assume that for some $k \in [1, \omega)$ the properties (I)–(VI) hold for $m = 1, \dots, k$. We again use Helly's theorem to obtain $p_{k+1} > p_k$ so that (V) holds for $m = k + 1$. To obtain that (VI) is true for $m = k + 1$ we modify the argument in the proof of (i) as follows. For $1 \leq i \leq k$ let $c_i \in \mathbb{R}$ be such that

$$k c_i \frac{4(2 + \nu)}{(1 - 2\nu)\epsilon_{n_i}} 2(1 + \nu) \sup\{\|\|y^*\| \mid y^* \in K\} = \frac{1}{2} \frac{\delta'_{k+1}}{3} \frac{(1 - 2\nu)\epsilon_{n_{k+1}}}{4(2 + \nu)} \frac{1}{(1 + \delta)^{1/2}}$$

and define

$$\mathcal{U}_3 = \bigcap_{i=1}^k \left\{ x^* \in X^* \mid |\langle x^* - f_{\tau(k+1)}^*, x_{\tau(i)} \rangle| < c_i \right\} \quad \text{and}$$

$$\mathcal{U}_4 = \bigcap_{p=1}^{p_k} \left\{ x^* \in X^* \mid |\langle x^* - f_{\tau(k+1)}^*, z_p \rangle| < \frac{1}{2} \frac{\delta'_{k+1}}{3} \frac{(1 - 2\nu)\epsilon_{n_{k+1}}}{4(2 + \nu)} \right\}.$$

We modify the definition of \mathcal{U} in the proof of (i) so that $\mathcal{U} := \mathcal{U}_1 \cap \mathcal{U}_2 \cap \mathcal{U}_3 \cap \mathcal{U}_4$ and, as in the proof of (i), choose $u^* \in \mathcal{U} \cap S_{\epsilon_{n_{k+1}}/2}^{\rho_{T_{n_{k+1}}}(t_{k+1})} \left((1 + \nu \sum_{j=1}^k 2^{-j}) K \right)$ such that $\|\|f_{\tau(k+1)}^* - u^*\| > \epsilon_{n_{k+1}}/4$. Now we define $f_{\tau(k+1)}^*$ and $x_{\tau(k+1)}$ as before. It is clear that properties (I)–(III) hold for them (by the same proof as before). Since (II) holds for $1 \leq m \leq k$ and since $u^* \in \mathcal{U}_3$, it follows from the definition of $f_{\tau(k+1)}^*$ that

$$\|\|f_{\tau(k+1)}^* - u^*\| \leq \frac{1}{2} \frac{\delta'_{k+1}}{3} \frac{(1 - 2\nu)\epsilon_{n_{k+1}}}{4(2 + \nu)} \frac{1}{(1 + \delta)^{1/2}}.$$

3. Absolutely convex sets of large Szlenk index

Hence, since $u^* \in \mathcal{U}_4$ we deduce that for all $p \leq p_k$ we have

$$\begin{aligned}
 & |\langle f_{\tau(k+1)}^* - f_{\tau(k+1)^-}^*, z_p \rangle| \\
 & \leq \|f_{\tau(k+1)}^* - u^*\| \|z_p\| + |\langle u^* - f_{\tau(k+1)^-}^*, z_p \rangle| \\
 & \leq \frac{1}{2} \frac{\delta'_{k+1}}{3} \frac{(1-2\nu)\epsilon_{n_{k+1}}}{4(2+\nu)} \frac{1}{(1+\delta)^{1/2}} (1+\delta)^{1/2} + \frac{1}{2} \frac{\delta'_{k+1}}{3} \frac{(1-2\nu)\epsilon_{n_{k+1}}}{4(2+\nu)} \\
 & = \frac{\delta'_{k+1}}{3} \frac{(1-2\nu)\epsilon_{n_{k+1}}}{4(2+\nu)}. \tag{18}
 \end{aligned}$$

Since

$$\begin{aligned}
 \|f_{\tau(k+1)}^* - f_{\tau(k+1)^-}^*\| & \geq \|f_{\tau(k+1)}^* - f_{\tau(k+1)^-}^*\| \\
 & \geq \langle f_{\tau(k+1)}^* - f_{\tau(k+1)^-}^*, x_{\tau(k+1)} \rangle > \frac{(1-2\nu)\epsilon_{n_{k+1}}}{4(2+\nu)},
 \end{aligned}$$

it follows from Equation (18) that for

$$v_{k+1} := \|f_{\tau(k+1)}^* - f_{\tau(k+1)^-}^*\|^{-1} (f_{\tau(k+1)}^* - f_{\tau(k+1)^-}^*)$$

we have $|\langle v_{k+1}, z_p \rangle| \leq \delta'_{k+1}/3$ for all $p \leq p_k$. Thus (I)–(VI) hold for all $m \in [1, \omega)$ with these modifications to the proof of (i).

Still following the argument in Johnson and Rosenthal (1972), our next step is to show that $(v_m)_{m=1}^\infty$ is a basic sequence whose basis constant with respect to $\|\cdot\|$ is no larger than $(1+\delta)^{1/2}$. To this end fix $m \in [1, \omega)$ and let $v \in [(v_q)_{q=1}^m]$ be such that $\|v\| = 1$. Choose $v^* \in [v_j]_{1 \leq j \leq m}^*$ such that $\langle v^*, v \rangle = 1 = \|v^*\|$ and choose $p \leq p_m$ so that (V) holds for v^* . Then $|\langle v, z_p \rangle| \geq 1 - \delta'_m/3$, hence for any scalar a we have

$$\begin{aligned}
 \|v + av_{m+1}\| & \geq \begin{cases} 1 & \text{if } |a| > 2 \\ |\langle v, z_p \rangle + \langle av_{m+1}, z_p \rangle| \geq (1 - \frac{\delta'_m}{3}) - \frac{2\delta'_m}{3} & \text{if } |a| \leq 2 \end{cases} \\
 & \geq 1 - \delta'_m.
 \end{aligned}$$

It follows that $\|\sum_{q=1}^m a_q v_q\| \leq \frac{1}{1-\delta'_m} \|\sum_{q=1}^{m+1} a_q v_q\|$ for any scalars a_1, \dots, a_m, a_{m+1} . Thus for $1 \leq l < m < \omega$ and any scalars a_1, \dots, a_m we have

$$\left\| \sum_{q=1}^l a_q v_q \right\| \leq \frac{1}{\prod_{q=l+1}^m (1-\delta'_q)} \left\| \sum_{q=1}^m a_q v_q \right\| \leq (1+\delta)^{1/2} \left\| \sum_{q=1}^m a_q v_q \right\|. \tag{19}$$

By Grunblum's criterion, $(v_m)_{m=1}^\infty$ is a basic sequence whose basis constant with respect to $\|\cdot\|$ is no larger than $(1+\delta)^{1/2}$.

Let $(v_m^*)_{m=1}^\infty$ be the sequence of functionals in $[(v_m)_{m=1}^\infty]^*$ biorthogonal to $(v_m)_{m=1}^\infty$ and define $T : X \rightarrow [(v_m)_{m=1}^\infty]^*$ by $\langle Tx, v \rangle = \langle v, x \rangle$ for $x \in X$ and $v \in [(v_m)_{m=1}^\infty]$. That

is, $Tx = (t_X x)|_{[(v_m)_{m=1}^\infty]}$ for each $x \in X$. Note that $\ker(T) = \bigcap_{m=1}^\infty \ker(v_m)$. Moreover since $f_{\tau(0)}^* = 0$ we have

$$f_t^* = \sum_{\emptyset < \mathfrak{I} \leq \mathfrak{I} t} (f_s^* - f_{s^-}^*)$$

for each $t \in \llbracket \mathfrak{I} \rrbracket$, hence

$$\ker(T) = \bigcap_{m=1}^\infty \ker(v_m) = \bigcap_{t \in \llbracket \mathfrak{I} \rrbracket^*} \ker(f_t^* - f_{t^-}^*) = \bigcap_{t \in \llbracket \mathfrak{I} \rrbracket^*} \ker(f_t^*) = \bigcap_{t \in \llbracket \mathfrak{I} \rrbracket^*} \ker(x_t^*) = Z.$$

It is a well known fact that the sequence of coordinate functionals associated with a basis is also a basis with the same basis constant³⁰, so let us momentarily denote by P_m , $m \geq 1$, the basis projections associated to the basic sequence $(v_m^*)_{m=1}^\infty$. Notice that by Equation (19) we have $\|P_m\| \rightarrow 1$ as $m \rightarrow \infty$. This together with the condition (V) is enough to prove that $T(X) = [(v_m^*)_{m=1}^\infty]$ (see pp. 83–84 in Johnson and Rosenthal 1972). Therefore there exists a linear isometry $\bar{T} : (X/Z, \|\cdot\|) \rightarrow ((v_m^*)_{m=1}^\infty, \|\cdot\|)$ such that $\bar{T}Qx = Tx$ for every $x \in X$. We thus have

$$\begin{aligned} \bar{T}Qx_{\tau(m)} &= Tx_{\tau(m)} = \sum_{m'=1}^\infty \langle Tx_{\tau(m)}, v_{m'} \rangle v_{m'}^* \\ &= \sum_{m'=1}^\infty \langle v_{m'}, x_{\tau(m)} \rangle v_{m'}^* = \frac{\langle f_{\tau(m)}^*, x_{\tau(m)} \rangle}{\|f_{\tau(m)}^* - f_{\tau(m)^-}^*\|} v_{m'}^*. \end{aligned}$$

For each $m \in [1, \omega)$ let $a_m = \langle f_{\tau(m)}^*, x_{\tau(m)} \rangle \|f_{\tau(m)}^* - f_{\tau(m)^-}^*\|^{-1}$, so that $\bar{T}Qx_{\tau(m)} = a_m v_m^*$ for each such m . As \bar{T} is an isometry with respect to $\|\cdot\|$, then, with respect to $\|\cdot\|$, $(Qx_{\tau(m)})_{m=1}^\infty$ is a basis for X/Z isometrically equivalent to $(a_m v_m^*)$, whose basis constant coincides with the basis constant of $(v_m^*)_{m=1}^\infty$, which coincides with the basis constant of $(v_m)_{m=1}^\infty$, which is no larger than $(1 + \delta)^{1/2}$ (as shown above). It follows that for $1 \leq l \leq m < \omega$ and any scalars a_1, \dots, a_m we have

$$\begin{aligned} \left\| \sum_{q=1}^l a_q Qx_{\tau(q)} \right\| &\leq (1 + \delta)^{1/2} \left\| \sum_{q=1}^l a_q Qx_{\tau(q)} \right\| \\ &\leq (1 + \delta) \left\| \sum_{q=1}^m a_q Qx_{\tau(q)} \right\| \\ &\leq (1 + \delta) \left\| \sum_{q=1}^m a_q Qx_{\tau(q)} \right\|. \end{aligned}$$

3. Absolutely convex sets of large Szlenk index

In particular, $(Qx_{\tau(m)})_{m=1}^{\infty}$ is a basis for X/Z whose basis constant with respect to $\|\cdot\|$ is no larger than $1 + \delta$. It remains then to show that $(Qx_{\tau(m)})_{m=1}^{\infty}$ is shrinking. As $(Qx_{\tau(m)})_{m=1}^{\infty}$ is equivalent to $(a_m v_m^*)_{m=1}^{\infty}$, which is shrinking if and only if $(v_m^*)_{m=1}^{\infty}$ is shrinking, it follows from the duality between shrinking and boundedly complete bases³¹ that, to complete the proof, it suffices to show that $(v_m)_{m=1}^{\infty}$ is boundedly complete. To this end recall the following definition from Johnson and Rosenthal (1972):

Definition 3 – Let X be a Banach space. A sequence $(y_m^*)_{m=1}^{\infty} \subseteq X^*$ is said to be *weak*-basic* if there is a sequence $(y_m)_{m=1}^{\infty} \subseteq X$ so that $(y_m, y_m^*)_{m=1}^{\infty}$ is biorthogonal and for each $y^* \in \overline{[y_m^* \mid m \in \mathbb{N}]}^{w^*}$ we have $\sum_{q=1}^m \langle y^*, y_q \rangle y_q^* \xrightarrow{\text{weak}^*} y^*$ as $m \rightarrow \infty$.

The following result is Proposition II.1 of Johnson and Rosenthal (1972).

Proposition 12 – Let X be a Banach space and $(y_m^*)_{m=1}^{\infty} \subseteq X^*$. Let Q denote the quotient map $X \rightarrow X / \bigcap_{m=1}^{\infty} \ker(y_m^*)$. Then

(a) $(y_m^*)_{m=1}^{\infty}$ is weak*-basic if and only if $X / \bigcap_{m=1}^{\infty} \ker(y_m^*)$ has a basis $(e_m)_{m=1}^{\infty}$ with associated biorthogonal functionals $(e_m^*)_{m=1}^{\infty}$ such that $Q^* e_m^* = y_m^*$ for all $m \in \mathbb{N}$. It follows that if $(y_m^*)_{m=1}^{\infty}$ is weak*-basic, then $(y_m^*)_{m=1}^{\infty}$ is basic.

(b) The following are equivalent:

(i) $(y_m^*)_{m=1}^{\infty}$ is a boundedly complete weak*-basic sequence;

(ii) $(y_m^*)_{m=1}^{\infty}$ is weak*-basic and $[(y_m^*)_{m=1}^{\infty}] = \overline{[y_m^* \mid m \in \mathbb{N}]}^{w^*}$.

We shall first apply (a) of Proposition 12 to show that $(v_m)_{m=1}^{\infty}$ is weak*-basic, then apply (b) of Proposition 12 to deduce that $(v_m)_{m=1}^{\infty}$ is boundedly complete, as desired. For $m \in [1, \omega)$ let

$$e_m := \frac{\|f_{\tau(m)}^* - f_{\tau(m)}^*\|}{\langle f_{\tau(m)}^*, x_{\tau(m)} \rangle} Qx_{\tau(m)},$$

so that $\overline{T}e_m = v_m^*$. For $m \in [1, \omega)$ let $v_m^{**} := (1_{[(v_q)_{q=1}^{\infty}]} v_m) |_{[(v_q^*)_{q=1}^{\infty}]}$ and $e_m^* := \overline{T}^* v_m^{**}$, so that $(v_m^{**})_{m=1}^{\infty}$ and $(e_m^*)_{m=1}^{\infty}$ are the sequences of biorthogonal functionals associated to the basic sequences $(v_m^*)_{m=1}^{\infty}$ and $(e_m)_{m=1}^{\infty}$, respectively. For $1 \leq m < \omega$ and $x \in X$ we have

$$\langle Q^* e_m^*, x \rangle = \langle e_m^*, Qx \rangle = \langle v_m^{**}, \overline{T}Qx \rangle = \langle Tx, v_m \rangle = \langle v_m, x \rangle, \quad (20)$$

hence $Q^* e_m^* = v_m$ for each $m \in [1, \omega)$. By Proposition 12(a), $(v_m)_{m=1}^{\infty}$ is weak*-basic. By Proposition 12(b), to complete the proof of Theorem 9 it now suffices to show that $[(v_m)_{m=1}^{\infty}] = \overline{[v_m \mid m \in \mathbb{N}]}^{w^*}$.

For each $m \in [1, \omega)$ let $y_m = \langle v_m, x_{\tau(m)} \rangle^{-1} x_{\tau(m)}$ so that, by Equation (9) and the definition of v_m , the system $(y_m, v_m)_{m=1}^{\infty} \subseteq X \times X^*$ is biorthogonal. Following now

the proof of Theorem III.2 in Johnson and Rosenthal (1972), for each $M \in [1, \omega)$ the operator

$$S_M : \overline{[v_m \mid m \in \mathbb{N}]^{w^*}} \longrightarrow \overline{[v_m \mid m \in \mathbb{N}]^{w^*}}$$

given by setting $S_M y^* = \sum_{m=1}^M \langle y^*, y_m \rangle v_m$ for each $y^* \in \overline{[v_m \mid m \in \mathbb{N}]^{w^*}}$ satisfies $\|S_M\| \leq \prod_{m=M}^{\infty} \frac{1}{1-\delta'_m}$. Suppose $y^* \in \overline{[v_m \mid m \in \mathbb{N}]^{w^*}}$. Then $(S_M y^*)_{M=1}^{\infty}$ converges weak* to y^* since $(v_m)_{m=1}^{\infty}$ is weak*-basic, hence $\liminf_M \|S_M y^*\| \geq \|y^*\|$. On the other hand, since $\|S_M\| \rightarrow 1$ we have $\limsup_M \|S_M y^*\| \leq \|y^*\|$. It follows that $\|S_M y^*\| \rightarrow \|y^*\|$ as $M \rightarrow \infty$, hence $\|S_M y^* - y^*\| \rightarrow 0$ since $\|\cdot\|$ satisfies property (ii) of Theorem 10. As $S_M y^* \in [(v_m)_{m=1}^{\infty}]$ for all M , we conclude that $y^* \in [(v_m)_{m=1}^{\infty}]$, which completes the proof of Theorem 9. \square

The following corollary of Theorem 9 may be useful in situation where one considers the ϵ -Szlenk index for only a single $\epsilon > 0$ (rather than a for a sequence $(\epsilon_n)_{n < \omega}$), such as the work in the current paper on universal operators.

Corollary 1 – *Let X be a Banach space, $K \subseteq X^*$ an absolutely convex, weak*-compact set, $\epsilon > 0$, $\xi > 0$ a countable ordinal, and (\mathcal{T}, \leq) a countable, well-founded tree such that $\rho(\mathcal{T}) \leq \xi + 1$. If $s_{\epsilon}^{\xi}(K) \neq \emptyset$ then there exist families $(x_t^*)_{t \in \mathcal{T}} \subseteq K$ and $(x_t)_{t \in \mathcal{T}} \subseteq S_X$ such that*

$$\langle x_t^*, x_s \rangle = \begin{cases} \langle x_s^*, x_s \rangle > \frac{\epsilon}{17} & \text{if } s \leq t \\ 0 & \text{if } s \not\leq t \end{cases}, \quad s, t \in \mathcal{T}. \quad (21)$$

Proof. Suppose $s_{\epsilon}^{\xi}(K) \neq \emptyset$ so that, by Lemma 1, $s_{\epsilon/2}^{\xi+1}(K) \supseteq s_{\epsilon/2}^{\xi 2}(K) \neq \emptyset$. Let t_0 be a set such that $t_0 \notin \mathcal{T}$ and let (\mathcal{T}_0, \leq_0) be the tree obtained by setting $\mathcal{T}_0 = \mathcal{T} \cup \{t_0\}$ and extending \leq to \mathcal{T}_0 by making t_0 the unique minimal element of \mathcal{T}_0 . Let $\xi_0 = \xi + 1$, so that $\rho(\mathcal{T}_0) \leq \xi_0 + 1$ and $s_{\epsilon/2}^{\xi_0}(K) \neq \emptyset$. The conclusion of the corollary follows from an application of Theorem 9(i) with $\theta = 1/2$, $\epsilon_n = \epsilon/2$ for all $n < \omega$, $\xi_n = 0$ for $0 < n < \omega$, and (\mathcal{T}_n, \leq_n) a tree consisting of a single node for $0 < n < \omega$ (since $([\mathcal{T}]^*, \leq_{\mathcal{T}}) = (\{0\} \times \mathcal{T}, \leq_{\mathcal{T}})$ is, in this case, naturally order isomorphic to \mathcal{T}). \square

4 Structure of Banach spaces of large Szlenk index

In this section we provide the proofs of Theorem 1 and Theorem 2. In doing so we continue with the notation introduced in Section 3. First we prove Theorem 1.

²⁹See e.g. Albiac and Kalton, 2016, *Topics in Banach space theory*, Proposition 1.1.9.

³⁰See e.g. Fabian et al., 2001, *Functional analysis and infinite-dimensional geometry*, Fact 6.6.

³¹See e.g. Singer, 1970, *Bases in Banach spaces. I*, Corollary 6.1.

4. Structure of Banach spaces of large Szlenk index

Proof (Proof of Theorem 1). Fix $\theta \in (0, \sqrt{65} - 8)$ and fix $\{\epsilon_n \mid n < \omega\}$, a countable dense subset of $(0, \infty) \subseteq \mathbb{R}$. We apply Theorem 9 with $K = B_{X^*}$, $\xi_n = \text{Sz}(X, \epsilon_n) - 1$ for each $n < \omega$, and $\mathfrak{T} = ((T_n, \leq_n))_{n < \omega}$ a family of blossomed trees with $\rho(T_n) = \text{Sz}(X, \epsilon_n)$ for each $n < \omega$, to obtain families $(x_t)_{t \in \llbracket \mathfrak{T} \rrbracket^*} \subseteq S_X$ and $(x_t^*)_{t \in \llbracket \mathfrak{T} \rrbracket^*} \subseteq B_{X^*}$ such that

$$\langle x_t^*, x_s \rangle = \begin{cases} \langle x_s^*, x_s \rangle > \frac{\epsilon_n}{8+\theta} & \text{if } s \leq_{\mathfrak{T}} t \in \{n\} \times T_n^* \\ 0 & \text{if } s \not\leq_{\mathfrak{T}} t \end{cases}, \quad s, t \in \llbracket \mathfrak{T} \rrbracket^*, n < \omega. \quad (22)$$

Let $Y_0 = [x_t]_{t \in \llbracket \mathfrak{T} \rrbracket^*}$. It follows from Equation (22) that for every $n < \omega$ the set $\{x_{(n,t)}^* \mid t \in T_n^*\}$ is $\frac{\epsilon_n}{8+\theta}$ -separated and that for every $t \in T_n^*$ the sequence $(x_{(n,u)}^*)_{u \in T_n[t+]}$ converges weak* in Y_0^* to $x_{n,t}^*$. Now an easy inductive argument (similar to that used to prove Equation (8)) shows that

$$\forall n < \omega \quad \forall (n, t) \in \{n\} \times T_n^* \quad x_{(n,t)}^*|_{Y_0} \in s_{\epsilon_n/(8+\theta)}^{\rho_{T_n^*}(t)}(B_{Y_0^*}).$$

Hence, since $(x_{(n,u)}^*)_{u \in T_n[\emptyset+]}$ converges weak* in Y_0^* to 0, we obtain that

$$0 \in s_{\epsilon_n/(8+\theta)}^{\rho(T_n)-1}(B_{Y_0^*}) = s_{\epsilon_n/(8+\theta)}^{\text{Sz}(X, \epsilon_n)-1}(B_{Y_0^*})$$

for each $n < \omega$. It follows that

$$\forall n < \omega \quad \text{Sz}\left(Y_0, \frac{\epsilon_n}{8+\theta}\right) \geq \text{Sz}(X, \epsilon_n). \quad (23)$$

Let $\tau : \omega \rightarrow \llbracket \mathfrak{T} \rrbracket$ be a bijection such that $\tau(m) \leq_{\mathfrak{T}} \tau(m')$ implies $m \leq m'$. Since $\text{Sz}(Y_0) < \omega_1$ and Y_0 is separable, the dual Y_0^* is norm separable. Thus, by Equation (23) and Theorem 9 there exist families $(y_t)_{t \in \llbracket \mathfrak{T} \rrbracket^*} \subseteq S_{Y_0}$ and $(y_t^*)_{t \in \llbracket \mathfrak{T} \rrbracket^*} \subseteq B_{Y_0^*}$ such that

$$\langle y_t^*, y_s \rangle = \begin{cases} \langle y_s^*, y_s \rangle > \frac{\epsilon_n}{65} & \text{if } s \leq_{\mathfrak{T}} t \in \{n\} \times T_n^* \\ 0 & \text{if } s \not\leq_{\mathfrak{T}} t \end{cases}, \quad s, t \in \llbracket \mathfrak{T} \rrbracket^*, n < \omega. \quad (24)$$

and $(y_{\tau(m)}^*)_{m=1}^{\infty}$ is a shrinking basic sequence with basis constant not exceeding $1 + \delta$. Let $Y = [(y_{\tau(m)}^*)_{m=1}^{\infty}]$. It follows from Equation (24) that

$$\forall n < \omega \quad \forall (n, t) \in \{n\} \times T_n^* \quad y_{(n,t)}^*|_Y \in s_{\epsilon_n/65}^{\rho_{T_n^*}(t)}(B_{Y^*})$$

and, subsequently, that $0 \in s_{\epsilon_n/65}^{\rho(T_n)-1}(B_{Y^*}) = s_{\epsilon_n/65}^{\text{Sz}(X, \epsilon_n)-1}(B_{Y^*})$ for each $n < \omega$. Thus,

$$\forall n < \omega \quad \text{Sz}\left(Y, \frac{\epsilon_n}{65}\right) \geq \text{Sz}(X, \epsilon_n). \quad (25)$$

For each $\epsilon > 0$ choose $N(\epsilon) < \omega$ such that $\epsilon_{N(\epsilon)} \in [\frac{65\epsilon}{66}, \epsilon]$. From Equation (25) we obtain

$$\forall \epsilon > 0 \quad \text{Sz}\left(Y, \frac{\epsilon}{66}\right) \geq \text{Sz}\left(Y, \frac{\epsilon_{N(\epsilon)}}{65}\right) \geq \text{Sz}(X, \epsilon_{N(\epsilon)}) \geq \text{Sz}(X, \epsilon),$$

which completes the proof of the theorem. \square

Two applications of Theorem 9 were used in the proof of Theorem 1 - the first to achieve separable reduction and the second to obtain a *shrinking* basic sequence. Clearly, if X is assumed norm separable then only one application of Theorem 9 is required, in which case the number 65 in Equation (1) may be replaced by $8 + \theta$ for any $\theta > 0$. Moreover, in the general case we may replace 65 by $16 + \theta$ for any $\theta > 0$; this is achieved by proving a version of Lemma 3.4 of Lancien (1996) for families $(\epsilon_n)_{n < \omega} \subseteq (0, \infty)$ and blossomed trees $((\mathcal{T}_n, \leq_n))_{n < \omega}$ (as in the proof Theorem 9), then applying this generalisation of Lemma 3.4 of Lancien (1996) to achieve separable reduction in the proof of Theorem 1 with $\epsilon/2$ (rather than $\epsilon/(8 + \theta)$) replacing ϵ .

We now prove Theorem 2.

Proof (Proof of Theorem 2). Fix a countable, dense subset $\{\epsilon_n \mid n < \omega\}$ of $(0, \infty)$. We apply Theorem 9 with $\theta = \frac{1}{2}$, $K = B_{X^*}$, $\xi_n = \text{Sz}(X, \epsilon_n) - 1$ for each $n < \omega$, and $\mathfrak{T} = ((\mathcal{T}_n, \leq_n))_{n < \omega}$ a family of blossomed trees with $\rho(\mathcal{T}_n) = \text{Sz}(X, \epsilon_n)$ for each $n < \omega$. Let $\tau : \omega \rightarrow \llbracket \mathfrak{T} \rrbracket^*$ be a bijection such that $\tau(m) \leq_{\mathfrak{T}} \tau(m')$ implies $m \leq m'$. By Theorem 9 there exist families $(x_t)_{t \in \llbracket \mathfrak{T} \rrbracket^*} \subseteq S_X$ and $(x_t^*)_{t \in \llbracket \mathfrak{T} \rrbracket^*} \subseteq B_{X^*}$ such that

$$\langle x_t^*, x_s \rangle = \begin{cases} \langle x_s^*, x_s \rangle > \frac{2\epsilon_n}{17} & \text{if } s \leq_{\mathfrak{T}} t \in \{n\} \times \mathcal{T}_n^* \\ 0 & \text{if } s \not\leq_{\mathfrak{T}} t \end{cases}, \quad s, t \in \llbracket \mathfrak{T} \rrbracket^*, n < \omega. \quad (26)$$

and $(Qx_{\tau(m)})_{m=1}^{\infty}$ is a shrinking basis for $X / \bigcap_{t \in \llbracket \mathfrak{T} \rrbracket^*} \ker(x_t^*)$ with basis constant not exceeding $1 + \delta$, where $Q : X \rightarrow X / \bigcap_{t \in \llbracket \mathfrak{T} \rrbracket^*} \ker(x_t^*)$ is the quotient map. Let $Z = \bigcap_{t \in \llbracket \mathfrak{T} \rrbracket^*} \ker(x_t^*)$. To complete the proof we will show that Equation (2) holds.

We may assume that the families $(x_t)_{t \in \llbracket \mathfrak{T} \rrbracket^*}$ and $(x_t^*)_{t \in \llbracket \mathfrak{T} \rrbracket^*}$ above are those constructed in the proof of Theorem 9. Let $(f_t^*)_{t \in \llbracket \mathfrak{T} \rrbracket^*}$ and $(v_m)_{m=1}^{\infty}$ also be as in the proof of Theorem 9. We have

$$\text{span}\{x_t^* \mid t \in \llbracket \mathfrak{T} \rrbracket^*\} = \text{span}\{f_t^* \mid t \in \llbracket \mathfrak{T} \rrbracket^*\} = \text{span}\{v_m \mid 1 \leq m < \omega\} \subseteq Q^*((X/Z)^*), \quad (27)$$

where the first equality is immediate from the definitions, the second equality follows from the inductively verified fact that

$$\forall k < \omega \quad \text{span}\{f_{\tau(m)}^* \mid 1 \leq m \leq k\} = \text{span}\{v_m \mid 1 \leq m \leq k\},$$

and the final inclusion follows from Equation (20). Since $\|Q\| = 1$ and since Q^* is an isometric embedding it follows respectively that $Qx_t \in B_{X/Z}$ and that $(Q^*)^{-1}(x_t^*)$ is

5. Universal operators of large Szlenk index

a well-defined element of $B_{(X/Z)^*}$ for every $t \in \llbracket \mathfrak{T} \rrbracket^*$. Since for $s, t \in \llbracket \mathfrak{T} \rrbracket^*$ and $n < \omega$ we have

$$\langle (Q^*)^{-1}(x_t^*), Qx_s \rangle = \langle x_t^*, x_s \rangle = \begin{cases} \langle x_s^*, x_s \rangle > \frac{2\epsilon_n}{17} & \text{if } s \leq_{\mathfrak{T}} t \in \{n\} \times \mathcal{T}_n^* \\ 0 & \text{if } s \not\leq_{\mathfrak{T}} t \end{cases}$$

and, since $\text{span}\{Qx_t \mid t \in \llbracket \mathfrak{T} \rrbracket^*\}$ is norm dense in X/Z , we have that the family $\{(Q^*)^{-1}(x_{(n,t)}^*) \mid t \in \mathcal{T}_n^*\}$ is $\frac{2\epsilon_n}{17}$ -separated and that for every $t \in \mathcal{T}_n$ the sequence $((Q^*)^{-1}(x_{(n,u)}^*))_{u \in \mathcal{T}_n[t+]}$ converges weak* in $(X/Z)^*$ to $(Q^*)^{-1}(x_{(n,t)}^*)$. Now an easy inductive argument (similar to that used to prove Equation (8)) yields

$$\forall n < \omega \quad \forall (n, t) \in \{n\} \times \mathcal{T}_n^* \quad (Q^*)^{-1}(x_{(n,t)}^*) \in s_{2\epsilon_n/17}^{\rho_{\mathcal{T}_n^*}(t)}(B_{(X/Z)^*})$$

and, subsequently, that $0 \in s_{2\epsilon_n/17}^{\rho(\mathcal{T}_n)-1}(B_{(X/Z)^*}) = s_{2\epsilon_n/17}^{\text{Sz}(X, \epsilon_n)-1}(B_{(X/Z)^*})$ for each $n < \omega$. Thus,

$$\forall n < \omega \quad \text{Sz}\left(X/Z, \frac{2\epsilon_n}{17}\right) \geq \text{Sz}(X, \epsilon_n). \quad (28)$$

For each $\epsilon > 0$ choose $N(\epsilon) < \omega$ such that $\epsilon_{N(\epsilon)} \in [\frac{17\epsilon}{18}, \epsilon]$. From Equation (28) we obtain

$$\forall \epsilon > 0 \quad \text{Sz}\left(X/Z, \frac{\epsilon}{9}\right) \geq \text{Sz}\left(X/Z, \frac{2\epsilon_{N(\epsilon)}}{17}\right) \geq \text{Sz}(X, \epsilon_{N(\epsilon)}) \geq \text{Sz}(X, \epsilon),$$

which completes the proof of the theorem. \square

5 Universal operators of large Szlenk index

The first and main undertaking of this section is the proof of Theorem 6 which provides the classification of the ordinals β for which the class $\mathcal{C}\mathcal{S}\mathcal{L}_\beta$ admits a universal element.

Proof (Proof of Theorem 6). Fix $\epsilon' > 0$ small enough that $s_{\epsilon'}^{\omega^\alpha}(T^*(B_{Y^*})) \neq \emptyset$ and fix $N < \omega$ large enough that $\rho(\mathcal{T}) \leq \omega^\alpha 2^N + 1$. Set $\epsilon = 2^{-N-1}\epsilon'$, so that $s_\epsilon^{\rho(\mathcal{T})-1}(T^*(B_{Y^*})) \neq \emptyset$ by Equation (10). Corollary 1 yields families $(x_t)_{t \in \mathcal{T}^*} \subseteq S_X$ and $(x_t^*)_{t \in \mathcal{T}^*} \subseteq T^*B_{Y^*}$ such that

$$\langle x_t^*, x_s \rangle = \begin{cases} \langle x_s^*, x_s \rangle > \frac{\epsilon}{17} & \text{if } s \leq t \\ 0 & \text{if } s \not\leq t' \end{cases}, \quad s, t \in \mathcal{T}^*.$$

By Proposition 8, $\Sigma_{\mathcal{T}^*}$ factors through T .

We now suppose that \mathcal{T} is blossomed and $\rho(\mathcal{T}) \geq \omega^\alpha$. Since \mathcal{T} is infinite and rooted we have $\rho(\mathcal{T}) \geq 2$ and that $\rho(\mathcal{T})$ is a successor ordinal. It follows that

$\rho(T) > \omega^\alpha$, hence by Proposition 11 we have $Sz(\Sigma_{T^\star}) \geq Sz(\Sigma_{T^\star}, \epsilon) \geq \rho(T) > \omega^\alpha$, so that $\Sigma_{T^\star} \in \mathcal{C}\mathcal{S}\mathcal{Z}_\alpha$. It follows that Σ_{T^\star} is universal for $\mathcal{C}\mathcal{S}\mathcal{Z}_\alpha$.

Finally, let β be an arbitrary ordinal. If $\beta < \omega_1$ then, by the second assertion of Theorem 6, Σ_{T^\star} is universal for $\mathcal{C}\mathcal{S}\mathcal{Z}_\beta$, where T_{ω^β} is as constructed in Example 2. Now suppose on the other hand that $\beta \geq \omega_1$; to complete the proof we show that $\mathcal{C}\mathcal{S}\mathcal{Z}_\beta$ does not admit a universal element. Suppose by way of contraposition that $\mathcal{C}\mathcal{S}\mathcal{Z}_\beta$ does admit a universal element, Υ say. Since³² $Sz(C(\omega^{\omega^\beta} + 1)) = \omega^{\beta+1}$, where $C(\omega^{\omega^\beta} + 1)$ denotes the Banach space of continuous scalar-valued functions on the compact ordinal $\omega^{\omega^\beta} + 1$, we have that Υ factors through the identity operator of $C(\omega^{\omega^\beta} + 1)$. It follows that $Sz(\Upsilon)$ satisfies $Sz(\Upsilon) \leq Sz(C(\omega^{\omega^\beta} + 1)) = \omega^{\beta+1}$ by the ideal property of $\mathcal{S}\mathcal{Z}_{\beta+1}$. Moreover the identity operator of ℓ_1 belongs to $\mathcal{C}\mathcal{S}\mathcal{Z}_\beta$ since ℓ_1 is not an Asplund space, hence Υ factors through ℓ_1 and, in particular, Υ has separable range. It thus follows by Proposition 4 that $Sz(\Upsilon) < \omega^{\omega_1} = \omega_1$, hence $\Upsilon \in \mathcal{S}\mathcal{Z}_{\omega_1} \subseteq \mathcal{S}\mathcal{Z}_\beta$ – a contradiction. Thus $\mathcal{C}\mathcal{S}\mathcal{Z}_\beta$ does not admit a universal element whenever $\beta \geq \omega_1$. \square

In Section 7 we will study whether each of the classes $\mathcal{D} \cap \mathcal{C}\mathcal{S}\mathcal{Z}_\beta$, $\beta \geq \omega_1$, admits a universal element of the form Σ_T for an uncountable tree T .

Remark 4 – It is straightforward to observe that we may replace Σ_{T^\star} by Σ_T in the statement of Theorem 6. However, the reason for our choice of Σ_{T^\star} over Σ_T is that universal operators may be thought of as ‘minimal’ elements of the class for which they are universal, and the operator Σ_{T^\star} can be thought of as naturally ‘smaller’ than Σ_T since T^\star is a subtree of T and Σ_{T^\star} therefore factors through Σ_T by Proposition 9. Moreover, T is not order isomorphic to a subtree of T^\star since T is assumed to be well-founded.

The following result can be proved directly using standard Szlenk index techniques, however as we shall refer to this result later we provide a quick proof here using Theorem 6.

Corollary 2 – *Let (T, \leq) be an infinite blossomed tree. Then $Sz(\Sigma_{T^\star}) = \rho(T)\omega$.*

Proof. Let $\alpha < \omega_1$ be the (unique) ordinal satisfying $\omega^\alpha \leq \rho(T) < \omega^{\alpha+1}$. Since $Sz(C(\omega^{\omega^\alpha} + 1)) = \omega^{\alpha+1} > \omega^\alpha$ by Samuel’s computation³³ of $Sz(C(K))$ for countable compact Hausdorff K , by Theorem 6 we have that Σ_{T^\star} factors through $C(\omega^{\omega^\alpha} + 1)$, hence $Sz(\Sigma_{T^\star}) \leq Sz(C(\omega^{\omega^\alpha} + 1)) = \omega^{\alpha+1} = \rho(T)\omega$ by the ideal property of $\mathcal{S}\mathcal{Z}_{\alpha+1}$.

As T is infinite and rooted we have $\rho(T) \geq 2$. Moreover, as noted in Section 2.2, $\rho(T)$ is a successor ordinal. It follows that $\rho(T) > \omega^\alpha$, hence $Sz(\Sigma_{T^\star}) \geq Sz(\Sigma_{T^\star}, \epsilon) \geq \rho(T) > \omega^\alpha$ by Proposition 11. As $Sz(\Sigma_{T^\star})$ is a power of ω , we deduce that $Sz(\Sigma_{T^\star}) \geq \omega^{\alpha+1} = \rho(T)\omega$, which completes the proof. \square

³²Brooker, 2013, “Szlenk and w^* -dentability indices of the Banach spaces $C([0, \alpha])$ ”, Theorem 2.6.

5. Universal operators of large Szlenk index

Remark 5 – Theorem 4 may be obtained as an immediate consequence of Theorem 6 and the fact that $\mathcal{K} = \mathcal{SZ}_0$ by Proposition 2.3 of Brooker (2012).

The following proposition relates some of the factorisation results of the current paper to known relationships between various closed operator ideals.

Proposition 13 – *Let I be a cofinal subset of ω_1 and for each ordinal $\xi \in I$ let \mathcal{T}_ξ be a blossomed tree with $\rho(\mathcal{T}_\xi) = \xi + 1$. For $T \in \mathcal{X}$, i.e. T with separable range, the following are equivalent:*

- (i) T factors $\Sigma_{\mathcal{T}_\xi^*}$ for every $\xi \in I$.
- (ii) T factors Σ_Ω , where Ω is the full countably branching tree of Example 1.
- (iii) T factors $\Sigma_{\mathcal{T}}$ for every countable tree \mathcal{T} with $ht(\mathcal{T}) \leq \omega$.
- (iv) $T \in \mathbb{C}\mathcal{D}$.
- (v) $T \in \mathbb{C}\mathcal{X}^*$.
- (vi) $T \in \mathbb{C}\mathcal{SZ}_{\omega_1}$.
- (vii) $T \in \mathbb{C}\bigcup_{\alpha \in \text{Ord}} \mathcal{SZ}_\alpha$.
- (viii) $T \in \mathbb{C}\bigcup_{\alpha < \omega_1} \mathcal{SZ}_\alpha$.

The proof of Proposition 13 relies on the following result of the author³⁴.

Theorem 11 – *Let X and Y be Banach spaces and $T \in \mathcal{L}(X, Y)$. Suppose that at least one of X and Y is separable and that $T \notin \mathcal{X}^*(X, Y)$. Then Σ_Ω factors through T .*

Proof (Proof of Proposition 13). The equivalence of assertions (iv)–(viii) follows immediately from Proposition 4. To complete the proof it thus suffices to show that (v) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (i) \Rightarrow (viii).

To see that (v) \Rightarrow (ii), let X and Y be Banach spaces and $T \in \mathcal{X}(X, Y) \setminus \mathcal{X}^*(X, Y)$. Let $\tilde{T} : X \rightarrow \overline{T(X)}$ be the separable codomain operator given by setting $\tilde{T}x = Tx$ for all $x \in X$. Since $\tilde{T} \notin \mathcal{X}^*$, by Theorem 11 there exist $U \in \mathcal{L}(\ell_1(\Omega), X)$ and $V \in \mathcal{L}(\overline{T(X)}, \ell_\infty(\Omega))$ such that $V\tilde{T}U = \Sigma_\Omega$. Since $\ell_\infty(\Omega)$ is injective³⁵ V admits a continuous linear extension $\tilde{V} \in \mathcal{L}(Y, \ell_\infty(\Omega))$, and for such \tilde{V} we have $\tilde{V}TU = \Sigma_\Omega$. Thus (v) \Rightarrow (ii).

The implication (ii) \Rightarrow (iii) follows from Proposition 9 and the fact that every countable tree \mathcal{T} with $ht(\mathcal{T}) \leq \omega$ is order isomorphic to a subtree of Ω , whilst (iii) \Rightarrow (i) is immediate from the fact that blossomed trees are by definition countable and well-founded.

Finally, the implication (i) \Rightarrow (viii) is a consequence of Proposition 11. □

³³Samuel, 1984, “Indice de Szlenk des $C(K)$ (K espace topologique compact dénombrable)”.

³⁴Brooker, 2017, “Non-Asplund Banach spaces and operators”, Theorem 5.5.

³⁵See Lindenstrauss and Tzafriri, 1977, *Classical Banach spaces. I*, p. 105.

6 When the codomain is separable

It is evident from the definition of the operator $\Sigma_{\mathcal{T}}$ associated to a tree (\mathcal{T}, \leq) that the range of $\Sigma_{\mathcal{T}}$ is contained in the closed linear span in $\ell_{\infty}(\mathcal{T})$ of the indicator functions $\chi_{\mathcal{T}[t \leq]}$, for $t \in \mathcal{T}$. For example, the range of the universal non-compact operator $\ell_1 \hookrightarrow \ell_{\infty}$ of Johnson (c.f. Theorem 4) is contained in the subspace c_0 of ℓ_{∞} , whilst the range of the Lindenstrauss-Pełczyński universal non-weakly compact summation operator from ℓ_1 to ℓ_{∞} (c.f. Theorem 3) is contained in the subspace c of ℓ_{∞} consisting of all convergent scalar sequences. In both these cases, the range is contained (up to isometric isomorphism) in a separable $C(K)$ space. In the corresponding papers of Johnson³⁶ and Lindenstrauss-Pełczyński³⁷ it is noted that stronger versions of the universal operator theorems presented there hold under restriction to the class of operators having separable codomain. More precisely, it is noted in Johnson (1971) that if $T : X \rightarrow Y$ is noncompact and Y is separable, then T factors the formal identity operator from ℓ_1 to c_0 . Moreover, in Lindenstrauss and Pełczyński (1968) it is noted that if $T : X \rightarrow Y$ is non-weakly compact and Y is separable, then T factors the summation operator from ℓ_1 to c defined by $(a_n)_{n=1}^{\infty} \mapsto (\sum_{i=1}^n a_i)_{i=1}^{\infty}$. In a similar vein, in the current section we show that for every $\alpha < \omega_1$ there exists an operator Υ_{α} from ℓ_1 into a separable $C(K)$ space with $Sz(\Upsilon_{\alpha}) > \omega^{\alpha}$ and such that Υ_{α} factors through any $T : X \rightarrow Y$ with Y separable and $Sz(T) \not\leq \omega^{\alpha}$.

Let (\mathcal{T}, \leq) be a rooted and chain-complete tree and let t_{\emptyset} denote the root of \mathcal{T} . By Theorem 8 the coarse wedge topology of \mathcal{T} is compact Hausdorff, thus for such \mathcal{T} we shall denote by $C(\mathcal{T})$ the Banach space (with the supremum norm) of coarse-wedge-continuous scalar-valued functions on \mathcal{T} . We denote by $C_0(\mathcal{T})$ the codimension-1 subspace $\{f \in C(\mathcal{T}) \mid f(t_{\emptyset}) = 0\}$ of $C(\mathcal{T})$. Note that if \mathcal{T} is countable then $C(\mathcal{T})$ and $C_0(\mathcal{T})$ are norm separable.

The following definition establishes the class of operators from which we shall draw our examples of universal non- α -Szlenk operators with separable codomain. We note that although the definition can be adapted to trees of arbitrarily large height, such generality is unnecessary for our purposes.

Definition 4 – Let (\mathcal{T}, \leq) be a rooted, well-founded tree. Define $\sigma_{\mathcal{T}} : \ell_1(\mathcal{T}) \rightarrow C(\mathcal{T})$ by

$$(\sigma_{\mathcal{T}}x)(t) = \sum_{s \leq t} x(s), \quad x \in \ell_1(\mathcal{T}), t \in \mathcal{T}.$$

That is, $\sigma_{\mathcal{T}}$ is the unique element of $\mathcal{L}(\ell_1(\mathcal{T}), C(\mathcal{T}))$ that maps each $e_t \in \ell_1(\mathcal{T})$ to $\chi_{\mathcal{T}[t \leq]} \in C(\mathcal{T})$. Similarly, define $\delta_{\mathcal{T}}$ to be the unique element of $\mathcal{L}(\ell_1(\mathcal{T}^*), C_0(\mathcal{T}))$ that maps each $e_t \in \ell_1(\mathcal{T}^*)$ to $\chi_{\mathcal{T}[t \leq]} \in C_0(\mathcal{T})$.

³⁶Johnson, 1971, “A universal non-compact operator”.

³⁷Lindenstrauss and Pełczyński, 1968, “Absolutely summing operators in \mathcal{L}_p -spaces and their applications”.

6. When the codomain is separable

Notice that Proposition 11 holds true with $\mathring{\sigma}_T$ in place of Σ_{T^*} . Indeed, since $C_0(T)$ naturally embeds linearly and isometrically into $\ell_\infty(T^*)$ via the restriction map $R \in \mathcal{L}(C_0(T), \ell_\infty(T^*))$, defined by setting $R(f) = f|_{T^*}$ for each $f \in C_0(T)$, and since $\Sigma_{T^*} = R\mathring{\sigma}_T$, we have $\mathring{\sigma}_T^*(B_{C_0(T)^*}) = \Sigma_{T^*}^*(B_{\ell_\infty(T^*)^*})$. We thus deduce that $Sz(\mathring{\sigma}_T) = Sz(\Sigma_{T^*})$ since the Szlenk indices of $\mathring{\sigma}_T$ and Σ_{T^*} are determined by the same subset of $\ell_1(T^*)^*$.

Similarly to the comments in Remark 4 regarding Theorem 6, we note that although $\mathring{\sigma}_T$ may be replaced by σ_T in the statement of Theorem 7, we present Theorem 7 as stated since $\mathring{\sigma}_T$ may be viewed as being naturally ‘smaller’ than σ_T .

A smallness condition of some kind on Y is necessary for the first assertion of Theorem 7 to hold in general. To see this, by Corollary 2 it is enough to observe that for a countably infinite, rooted, well-founded tree (T, \leq) , $\mathring{\sigma}_T$ does not factor through Σ_{T^*} . Firstly, the fact that such T is infinite and well-founded implies that $\text{MAX}(T)$ contains an infinite anti-chain $\{t_n \mid n < \omega\}$, so that $\mathring{\sigma}_T$ is non-compact since the set $\{\mathring{\sigma}_T e_{t_n} \mid n < \omega\}$ is an infinite ϵ -separated subset of $\mathring{\sigma}_T(B_{\ell_1(T^*)})$ for any $\epsilon \in (0, 1)$. Secondly, the norm separability of $C_0(T)$ and the fact that $\ell_\infty(T^*)$ is a Grothendieck space implies that³⁸ $\mathcal{L}(\ell_\infty(T^*), C_0(T)) = \mathcal{W}(\ell_\infty(T^*), C_0(T))$. Thirdly, the fact that T is well-founded implies that Σ_{T^*} is weakly compact by Proposition 10, hence for any $V \in \mathcal{L}(\ell_\infty(T^*), C_0(T)) = \mathcal{W}(\ell_\infty(T^*), C_0(T))$ we have that $V\Sigma_{T^*}$ is compact since $\ell_\infty(T^*)$ is isomorphic to a $C(K)$ space and therefore has the Dunford-Pettis Property³⁹⁴⁰. Finally, since $V\Sigma_{T^*}$ therefore cannot factor $\mathring{\sigma}_T$ for any $V \in \mathcal{L}(\ell_\infty(T^*), C_0(T))$, we conclude that Σ_{T^*} does not factor $\mathring{\sigma}_T$.

To prove Theorem 7 we first establish the following continuous analogue of Proposition 8.

Proposition 14 – *Let K be a compact Hausdorff space, I an index set and $\{K_i\}_{i \in I}$ a family of clopen subsets of K . For Banach spaces X and Y and $T \in \mathcal{L}(X, Y)$ the following are equivalent:*

- (i) T factors the unique element of $\mathcal{L}(\ell_1(I), C(K))$ satisfying $e_i \mapsto \chi_{K_i}$, $i \in I$.
- (ii) There exists $(\delta_i)_{i \in I} \subseteq \mathbb{R}$ with $\inf_{i \in I} \delta_i > 0$, a family $(x_i)_{i \in I} \subseteq X$ with $\sup_{i \in I} \|x_i\| < \infty$ and a weak*-continuous map $\Xi : K \rightarrow Y^*$ such that

$$\forall i \in I \quad \forall k \in K \quad \langle \Xi(k), Tx_i \rangle = \begin{cases} \delta_i, & k \in K_i \\ 0, & k \notin K_i \end{cases}.$$

Proof. For each $k \in K$ let $g_k^* \in C(K)^*$ denote the evaluation functional of $C(K)$ at k ; that is, $g_k^*(f) = f(k)$ for each $f \in C(K)$.

³⁸Grothendieck, 1953, “Sur les applications linéaires faiblement compactes d’espaces du type $C(K)$ ”.

³⁹Ibid.

⁴⁰See also Albiac and Kalton, 2016, *Topics in Banach space theory*, Theorem 5.4.6.

Suppose (i) holds. Let S denote the unique element of $\mathcal{L}(\ell_1(I), C(K))$ satisfying $e_i \mapsto \chi_{K_i}$, $i \in I$, and let $U \in \mathcal{L}(\ell_1(I), X)$ and $V \in \mathcal{L}(Y, C(K))$ be such that $S = VTU$. The map $k \mapsto g_k^*$ is a homeomorphic embedding of K into $C(K)^*$ with respect to the weak*-topology of $C(K)^*$, hence the map $\Xi : K \rightarrow Y^*$ defined by setting $\Xi(k) = V^*g_k^*$ for each $k \in K$ is weak*-continuous. For each $i \in I$ set $x_i = Ue_i$, so that $\sup_{i \in I} \|x_i\| \leq \|U\| < \infty$. Then for $i \in I$ and $k \in K$ we have

$$\langle \Xi(k), Tx_i \rangle = \langle V^*g_k^*, T U e_i \rangle = \langle g_k^*, VT U e_i \rangle = \langle g_k^*, S e_i \rangle = \langle g_k^*, \chi_{K_i} \rangle = \begin{cases} 1, & k \in K_i \\ 0, & k \notin K_i \end{cases}.$$

By taking $\delta_i = 1$ for each $i \in I$ we see that (ii) holds, as desired.

Now suppose (ii) holds. Let $U \in \mathcal{L}(\ell_1(I), X)$ be defined by setting $Ue_i = \delta_i^{-1}x_i$ for each $i \in I$, noting that U is well-defined with $\|U\| \leq (\inf_{i \in I} \delta_i)^{-1} \sup_{i \in I} \|x_i\|$. Let V be the element of $\mathcal{L}(Y, C(K))$ satisfying $(Vy)(k) = \langle \Xi(k), y \rangle$ for $y \in Y$ and $k \in K$, noting that V is well-defined with $\|V\| = \sup_{k \in K} \|\Xi(k)\| < \infty$. For $i \in I$ and $k \in K$ we have

$$(VTUe_i)(k) = \langle g_k^*, VTUe_i \rangle = \delta_i^{-1} \langle V^*g_k^*, Tx_i \rangle = \delta_i^{-1} \langle \Xi(k), Tx_i \rangle = \begin{cases} 1, & k \in K_i \\ 0, & k \notin K_i \end{cases},$$

hence $VTUe_i = Se_i$ for every $i \in I$, hence $VTU = S$. \square

The following lemma is another key ingredient required for the proof of Theorem 7.

Lemma 5 – *Let X and Y be Banach spaces such that Y is separable. Let $T \in \mathcal{L}(X, Y)$, $\delta > 0$, (\mathcal{R}, \leq') a blossomed tree and $(x_r)_{r \in \mathcal{R}} \subseteq S_X$ and $(x_r^*)_{r \in \mathcal{R}} \subseteq T^*(B_{Y^*})$ families such that*

$$\langle x_s^*, x_r \rangle = \begin{cases} \langle x_r^*, x_r \rangle > \delta & \text{if } r \leq' s \\ 0 & \text{if } r \not\leq' s \end{cases}, \quad r, s \in \mathcal{R}. \quad (29)$$

Then for any $\xi < \rho(\mathcal{R})$ and any $r_0 \in \mathcal{R}^{(\xi)} \setminus \mathcal{R}^{(\xi+1)}$ there exists a full subtree \mathcal{S} of $\mathcal{R}[r_0 \leq']$ and a family $(y_s^)_{s \in \mathcal{S}} \subseteq B_{Y^*}$ such that*

$$\langle y_s^*, Tx_r \rangle = \langle x_s^*, x_r \rangle, \quad s \in \mathcal{S}, r \in \mathcal{R} \quad (30)$$

and the map $s \mapsto y_s^$ from \mathcal{S} to Y^* is coarse-wedge-to-weak* continuous.*

Proof. We proceed by induction on ξ . For the base case, namely $\xi = 0$, fix $r_0 \in \mathcal{R}^{(0)} \setminus \mathcal{R}^{(1)}$, let $\mathcal{S} = \{r_0\}$ and choose $y_{r_0}^* \in B_{Y^*}$ such that $T^*y_{r_0}^* = x_{r_0}^*$. In this way we see that the assertion of the lemma is true in the case $\xi = 0$.

We now address the inductive step. Suppose $0 < \zeta < \rho(\mathcal{R})$ and that the assertion of the lemma holds for all $\xi < \zeta$; we will now show it is then true for $\xi = \zeta$. To this

6. When the codomain is separable

end fix $r_0 \in \mathcal{R}^{(\zeta)} \setminus \mathcal{R}^{(\zeta+1)}$ and for each $t \in \mathcal{R}[r_0+]$ let \mathcal{S}_t be a full subtree of $\mathcal{R}[t \leq']$ and $(y_s^*)_{s \in \mathcal{S}_t} \subseteq B_{Y^*}$ a family such that

$$\langle y_s^*, Tx_r \rangle = \langle x_s^*, x_r \rangle, \quad s \in \mathcal{S}_t, r \in \mathcal{R} \quad (31)$$

and the map $\Xi_t : s \mapsto y_s^*$ from \mathcal{S}_t to Y^* is coarse-wedge-to-weak* continuous. Let d be a metric on B_{Y^*} that is compatible with the weak* topology on B_{Y^*} and let $(t_n)_{n=0}^\infty$ be an injective sequence in $\mathcal{R}[r_0+]$ such that $(y_{t_n}^*)_{n=0}^\infty$ is weak* convergent. Let $y_{r_0}^*$ denote the weak*-limit of $(y_{t_n}^*)_{n=0}^\infty$. Passing to a subsequence we may assume that $d(y_{t_n}^*, y_{r_0}^*) < 1/n$ for each $n < \omega$. By Proposition 6 and the continuity of the maps Ξ_{t_n} , $n < \omega$, for each $n < \omega$ we may choose a finite set $\mathcal{F}_n \subseteq \mathcal{S}_{t_n}[t_n+]$ such that

$$\forall n < \omega \quad \forall t \in \mathcal{W}_{\mathcal{S}_{t_n}}(t_n, \mathcal{F}_n) \quad d(\Xi_{t_n}(t), \Xi_{t_n}(t_n)) < \frac{1}{n}.$$

Define $\mathcal{S} := \{r_0\} \cup \bigcup_{n < \omega} \mathcal{W}_{\mathcal{S}_{t_n}}(t_n, \mathcal{F}_n)$ and $\Xi : \mathcal{S} \rightarrow Y^*$ by

$$\Xi(s) = \begin{cases} y_{r_0}^*, & s = r_0 \\ \Xi_{t_n}(s), & s \in \mathcal{W}_{\mathcal{S}_{t_n}}(t_n, \mathcal{F}_n), n < \omega \end{cases}$$

It is straightforward to check that \mathcal{S} is a full subtree of $\mathcal{R}[r_0 \leq']$ since $\mathcal{W}_{\mathcal{S}_{t_n}}(t_n, \mathcal{F}_n)$ is a full subtree of \mathcal{S}_{t_n} for every $n < \omega$. To see that Equation (30) holds for this \mathcal{S} , note that since Equation (31) holds for $t = t_n$, for all $n < \omega$, we need only check that Equation (30) holds in the case where $s = r_0$. To this end note that for all $r \in \mathcal{R}$ we have

$$\langle y_{r_0}^*, Tx_r \rangle = \lim_{n \rightarrow \omega} \langle y_{t_n}^*, Tx_r \rangle = \lim_{n \rightarrow \omega} \langle x_{t_n}^*, x_r \rangle = \langle x_{r_0}^*, x_r \rangle,$$

where the final equality follows from Equation (29). Thus Equation (30) holds for all $s \in \mathcal{S}$ and $r \in \mathcal{R}$.

To complete the proof it remains only to establish the continuity of Ξ . Since each Ξ_{t_n} is continuous, for $n < \omega$, the only nontrivial case to check is whether Ξ is continuous at r_0 . Fix $\lambda > 0$. Let $N < \omega$ be large enough that $N\lambda > 2$ and let $\mathcal{F} = \{t_0, \dots, t_{N-1}\}$. For each $s \in \mathcal{W}_{\mathcal{S}}(r_0, \mathcal{F}) \setminus \{r_0\}$ there exists a unique $n_s \geq N$ such that $t_{n_s} \leq' s$. So for $s \in \mathcal{W}_{\mathcal{S}}(r_0, \mathcal{F}) \setminus \{r_0\}$ we have

$$d(\Xi(s), \Xi(r_0)) \leq d(\Xi(s), \Xi(t_{n_s})) + d(\Xi(t_{n_s}), \Xi(r_0)) < \frac{1}{n_s} + \frac{1}{n_s} \leq \frac{2}{N} < \lambda.$$

It follows that $d(\Xi(s), \Xi(r_0)) < \lambda$ for all $s \in \mathcal{W}_{\mathcal{S}}(r_0, \mathcal{F})$, hence Ξ is continuous at r_0 since $\mathcal{W}_{\mathcal{S}}(r_0, \mathcal{F})$ is open in \mathcal{S} by Proposition 6. \square

Proof (Proof of Theorem 7). Suppose Y is separable. Let (\mathcal{R}, \leq') be a blossomed tree with $\rho(\mathcal{R}) = \rho(T)$ (c.f. Example 2), let $\epsilon' > 0$ be small enough that $s_{\epsilon'}^{\omega_\alpha}(T^*(B_{Y^*})) \neq \emptyset$,

and let $N < \omega$ be large enough $\rho(T) \leq \omega^\alpha 2^N + 1$. Set $\epsilon = 2^{-N-1}\epsilon'$, so that $s_\epsilon^{\rho(T)-1}(T^*(B_{Y^*})) \neq \emptyset$ by Equation (10). Corollary 1 yields families $(x_s)_{s \in \mathcal{R}} \subseteq S_X$ and $(x_s^*)_{s \in \mathcal{R}} \subseteq T^*(B_{Y^*})$ such that

$$\langle x_s^*, x_r \rangle = \begin{cases} \langle x_r^*, x_r \rangle > \frac{\epsilon}{17} & \text{if } r \leq' s \\ 0 & \text{if } r \not\leq' s \end{cases}, \quad r, s \in \mathcal{R}. \quad (32)$$

We apply Lemma 5 with r_0 the root of \mathcal{R} and $\xi = \rho(T) - 1$ to obtain a full subtree \mathcal{S} of \mathcal{R} and a family $(y_s^*)_{s \in \mathcal{S}} \subseteq B_{Y^*}$ such that

$$\langle y_s^*, Tx_r \rangle = \langle x_s^*, x_r \rangle, \quad s, r \in \mathcal{S}, \quad (33)$$

and the map $\Xi : s \mapsto y_s^*$ from \mathcal{S} to Y^* is coarse-wedge-to-weak* continuous. From Equation (32) and Equation (33) we deduce that

$$\langle y_s^*, Tx_r \rangle = \begin{cases} \langle y_r^*, Tx_r \rangle > \frac{\epsilon}{17} & \text{if } r \leq' s \\ 0 & \text{if } r \not\leq' s \end{cases}, \quad r, s \in \mathcal{S}. \quad (34)$$

By an application of Proposition 14 with $K = \mathcal{S}$, index set $I = \mathcal{S}$, clopen sets $K_s = \mathcal{S}[s \leq']$ for $s \in \mathcal{S}$, and $\delta_s = \epsilon/17$ for all $s \in \mathcal{S}$, we obtain that $\sigma_{\mathcal{S}}$ factors through T . So to prove the first assertion of Theorem 7 it now suffices to show that $\mathring{\sigma}_{\mathcal{T}}$ factors through $\sigma_{\mathcal{S}}$. To this end we now define three operators, S , R , and P , so that $\mathring{\sigma}_{\mathcal{T}} = PS\sigma_{\mathcal{S}}R$. Let $\phi : \mathcal{T} \rightarrow \mathcal{S}$ be an order-isomorphism of \mathcal{T} onto a downward-closed subtree of \mathcal{S} , noting that such an embedding exists by Proposition 5. Since ϕ is coarse wedge continuous by Proposition 7(ii), the operator $S \in \mathcal{L}(C(\mathcal{S}), C(\mathcal{T}))$ given by setting $Sf = f \circ \phi$ for each $f \in C(\mathcal{S})$ is well-defined. Let $R \in \mathcal{L}(\ell_1(T^*), \ell_1(\mathcal{S}))$ be the operator defined by setting $Re_t = e_{\phi(t)}^{\mathcal{S}}$ for every $t \in T^*$. Let t_{\emptyset} denote the root of \mathcal{T} and define $P \in \mathcal{L}(C(\mathcal{T}), C_0(\mathcal{T}))$ by setting $Pf = f - f(t_{\emptyset})\chi_{\mathcal{T}}$ for each $f \in C(\mathcal{T})$. Since for $t \in T^*$ we have

$$PS\sigma_{\mathcal{S}}Re_t = PS\sigma_{\mathcal{S}}e_{\phi(t)}^{\mathcal{S}} = PS\chi_{\mathcal{S}[\phi(t) \leq']} = P\chi_{\mathcal{T}[t \leq]} = \chi_{\mathcal{T}[t \leq]} = \mathring{\sigma}_{\mathcal{T}}e_t,$$

we conclude that $\mathring{\sigma}_{\mathcal{T}} = PS\sigma_{\mathcal{S}}R$. The first assertion of the theorem is proved.

For the second assertion of the theorem, we now suppose that \mathcal{T} is blossomed and $\rho(T) \geq \omega^\alpha$. As \mathcal{T} is infinite and rooted, we have that $\rho(T) \geq 2$ and $\rho(T)$ is a successor ordinal, hence $\rho(T) > \omega^\alpha$. Since \mathcal{T} is blossomed, an application of Proposition 11 yields $\text{Sz}(\Sigma_{\mathcal{T}^*}) \geq \rho(T) > \omega^\alpha$. Moreover, as noted in the paragraph following Definition 4 on p. 198, $\text{Sz}(\Sigma_{\mathcal{T}^*})$ coincides with $\text{Sz}(\mathring{\sigma}_{\mathcal{T}})$, hence $\mathring{\sigma}_{\mathcal{T}}$ is non- α -Szlenk. Note also that the codomain of $\mathring{\sigma}_{\mathcal{T}}$, namely $C_0(\mathcal{T})$, is norm separable since \mathcal{T} is countable. On the other hand, by the first assertion of the theorem we have that $\mathring{\sigma}_{\mathcal{T}}$ factors through any non- α -Szlenk operator with separable codomain, hence we conclude that $\mathring{\sigma}_{\mathcal{T}}$ is in this case universal for the class of non- α -Szlenk operators with separable codomain. \square

6. When the codomain is separable

Remark 6 – Bourgain, in a study⁴¹ of fixing properties of operators of large Szlenk index acting on $C(K)$ spaces, represented $C(L)$ spaces with L countable, compact and Hausdorff as spaces of scalar-valued functions on blossomed trees. Bourgain associates to each tree $(\mathcal{T}_\xi, \sqsubseteq)$ constructed in Example 2 above (subtrees of the tree (Ω, \sqsubseteq) constructed in Example 1) a Banach space X_ξ , isometrically isomorphic to $C_0(\mathcal{T}_\xi)$, defined as the completion of $c_{00}(\mathcal{T}_\xi^*)$ (the space of finitely-supported scalar-valued functions on \mathcal{T}_ξ^*) with respect to the norm $\|\cdot\|_\xi$ defined by setting

$$\|x\|_\xi = \sup_{t \in \mathcal{T}_\xi^*} \left| \sum_{s \sqsubseteq t} x(s) \right|, \quad x \in c_{00}(\mathcal{T}_\xi^*).$$

The main assertion of Theorem 7 may be recast as follows: *Suppose $\alpha, \xi < \omega_1$ are such that $\omega^\alpha \leq \xi < \omega^{\alpha+1}$ and let $T_\xi \in \mathcal{L}(\ell_1(\mathcal{T}_\xi^*), X_{\mathcal{T}_\xi^*})$ denote the continuous linear extension of the formal identity map from $(c_{00}(\mathcal{T}_\xi^*), \|\cdot\|_{\ell_1(\mathcal{T}_\xi^*)})$ to $X_{\mathcal{T}_\xi^*}$. Then T_ξ is universal for the class of non- α -Szlenk operators with norm separable codomain.*

We conclude the current section with some observations regarding the aforementioned universal operator theorems of Johnson and Lindenstrauss-Pełczyński. In particular, we note the following corollaries of Theorems 3 and 4. These results appear in Johnson (1971) and Lindenstrauss and Pełczyński (1968), respectively, under the stronger hypothesis that Y is norm separable.

Corollary 3 – *Let X and Y be Banach spaces such that Y has weak*-sequentially compact dual ball and let $T \in \mathcal{L}(X, Y)$ be non-weakly compact. Then T factors the summation operator $(a_n)_{n=1}^\infty \mapsto (\sum_{i=1}^n a_i)_{n=1}^\infty$ from ℓ_1 to c .*

Proof. By Theorem 3 there exist $U \in \mathcal{L}(\ell_1, X)$ and $V \in \mathcal{L}(Y, \ell_\infty)$ such that VTU is the summation operator from ℓ_1 to ℓ_∞ . For $n \in \mathbb{N}$ let f_n^* denote the n th coordinate functional on ℓ_∞ ; that is, $f_n^*(f) = f(n)$ for every $f \in \ell_\infty$. Since Y has weak*-sequentially compact dual ball there is a weak*-convergent subsequence $(V^* f_{n_k}^*)_{k=1}^\infty$ of $(V^* f_n^*)_{n=1}^\infty$. Define $A \in \mathcal{L}(\ell_1, X)$ by setting $Ae_k = Ue_{n_k}$ for each $k \in \mathbb{N}$ and define $B \in \mathcal{L}(Y, c)$ by setting $By = (\langle V^* f_{n_k}^*, y \rangle)_{k=1}^\infty$ for each $y \in Y$. For $k, l \in \mathbb{N}$ we have

$$(BTAe_k)(l) = \langle V^* f_{n_l}^*, TAe_k \rangle = \langle f_{n_l}^*, VTUe_{n_k} \rangle = \begin{cases} 1, & l \geq k \\ 0, & l < k \end{cases},$$

hence BTA coincides with the summation operator from ℓ_1 to c . □

Corollary 4 – *Let X and Y be Banach spaces such that Y has weak*-sequentially compact dual ball and let $T \in \mathcal{L}(X, Y)$ be non-compact. Then T factors the formal identity mapping from ℓ_1 to c_0 .*

⁴¹Bourgain, 1979, “The Szlenk index and operators on $C(K)$ -spaces”.

Proof. By Theorem 4 there exist $U \in \mathcal{L}(\ell_1, X)$ and $V \in \mathcal{L}(Y, \ell_\infty)$ such that VTU is the formal identity mapping from ℓ_1 to ℓ_∞ . For $n \in \mathbb{N}$ let f_n^* denote the n th coordinate functional on ℓ_∞ . Since Y has weak*-sequentially compact dual ball there is a subsequence $(V^*f_{n_k}^*)_{k=1}^\infty$ of $(V^*f_n^*)_{n=1}^\infty$ converging weak* to some $y^* \in Y^*$. Define $A \in \mathcal{L}(\ell_1, X)$ by setting $Ae_k = Ue_{n_k}$ for each $k \in \mathbb{N}$ and define $B \in \mathcal{L}(Y, c_0)$ by setting $By = (\langle V^*f_{n_k}^* - y^*, y \rangle)_{k=1}^\infty$ for each $y \in Y$. Since

$$\forall k \in \mathbb{N} \quad \langle y^*, T U e_{n_k} \rangle = \lim_{l \rightarrow \infty} \langle V^*f_{n_l}^*, T U e_{n_k} \rangle = \lim_{l \rightarrow \infty} \langle f_{n_l}^*, V T U e_{n_k} \rangle = 0,$$

it follows that for $k, l \in \mathbb{N}$ we have

$$(BTAe_k)(l) = (BT U e_{n_k})(l) = \langle V^*f_{n_l}^* - y^*, T U e_{n_k} \rangle = \langle f_{n_l}^*, V T U e_{n_k} \rangle = \begin{cases} 1, & l = k \\ 0, & l \neq k \end{cases},$$

hence BTA coincides with the formal identity mapping from ℓ_1 to c_0 . \square

We conclude the current section of the paper with the following open question.

Question 1 – Does the statement of Theorem 7 remain true if the condition that Y be norm separable is relaxed and Y is only assumed to have weak*-sequentially compact dual ball?

7 Uncountable Szlenk indices and universality

The proof of Theorem 6 makes use of the fact that an operator T can fail to be α -Szlenk for a given ordinal α in essentially two different ways. More precisely, it can be that $\text{Sz}(T)$ is defined and larger than ω^α , or it can be that $\text{Sz}(T)$ is undefined. In particular, a non α -Szlenk operator can be either Asplund or non-Asplund. This observation and the first assertion of Theorem 6 lead naturally to the following open question.

Question 2 – Let $\alpha \geq \omega_1$ be an uncountable ordinal. Does $\mathcal{D} \cap \mathcal{C}\mathcal{SZ}_\alpha$ admit a universal element?

In light of the approach taken in Theorem 6 to prove the existence of universal elements of $\mathcal{C}\mathcal{SZ}_\alpha$ for $\alpha < \omega_1$, it is natural to guess that a first step towards answering Question 2 could involve consideration of operators of the form Σ_T , where (T, \leq) is a tree, and extending the definition of a blossomed tree (c.f. Definition 2) to the uncountable setting as follows: say that a tree (T, \leq) is *blossomed* if it is rooted, well-founded, and for every $t \in T \setminus \text{MAX}(T)$ there exists a bijection $\psi_t : \max\{\omega, \text{cof}(\rho_T(t))\} \rightarrow T[t+]$ such that $\zeta \leq \zeta' < \max\{\omega, \text{cof}(\rho_T(t))\}$ implies $\rho_T(\psi_t(\zeta)) \leq \rho_T(\psi_t(\zeta'))$. Examples of such trees T with $\rho(T) = \xi + 1$ for a given ordinal ξ may be obtained via a similar construction to that provided in Example 2, but with T consisting of finite sequences of ordinals (ordered by extension, as in

7. Uncountable Szlenk indices and universality

Example 2). Moreover, under this more general definition of a blossomed tree, a blossomed tree (\mathcal{T}, \leq) satisfies $\rho(\mathcal{T}) < \omega_1$ if and only if \mathcal{T} is countable and satisfies the usual definition of blossomed tree given in Definition 2. The natural candidate for a universal element of $\mathcal{D} \cap \bigcap \mathcal{SZ}_{\omega_1}$ under this approach is $\Sigma_{\mathcal{S}}$, where for each $\alpha < \omega_1$ we let $(\mathcal{T}_\alpha, \leq_\alpha)$ be a blossomed tree with $\rho(\mathcal{T}_\alpha) = \alpha + 1$ and set $\mathcal{S} = \bigcup_{\alpha < \omega_1} (\{\alpha\} \times \mathcal{T}_\alpha)$, with an order \leq on \mathcal{S} defined by setting $(\alpha, t) \leq (\alpha', t')$ if and only if $\alpha = \alpha'$ and $t \leq_\alpha t'$. We do not know the answer to Question 2, even in the case $\alpha = \omega_1$. However, as we shall now see, operators of the form $\Sigma_{\mathcal{T}}$, where (\mathcal{T}, \leq) is a well-founded tree - cannot be expected to provide absolute examples of universal elements of the classes $\mathcal{D} \cap \bigcap \mathcal{SZ}_\alpha$ for $\alpha \geq \omega_1$ in general. In particular, we shall see that it is consistent with ZFC that the operator $\Sigma_{\mathcal{S}}$ defined above is not universal for $\mathcal{D} \cap \bigcap \mathcal{SZ}_{\omega_1}$.

Let Z be a Banach space and (\mathcal{T}, \leq) a tree. Let

$$\mathcal{O}_{\mathcal{T}} = \{t \in \mathcal{T} \mid ht_{\mathcal{T}}(t) = 0 \text{ or } ht_{\mathcal{T}}(t) \text{ is a successor}\},$$

noting that $\mathcal{O}_{\mathcal{T}} = \mathcal{T}$ if and only if $ht(\mathcal{T}) \leq \omega$. In particular, $\mathcal{O}_{\mathcal{T}} = \mathcal{T}$ whenever \mathcal{T} is well-founded. If $\Sigma_{\mathcal{T}}$ factors through Z then, by Proposition 8, there exist $\delta > 0$, $(x_t)_{t \in \mathcal{T}} \subseteq Z$ and $(x_t^*)_{t \in \mathcal{T}} \subseteq Z^*$ satisfying Equation (7). It follows that Z admits a biorthogonal system of cardinality $|\mathcal{O}_{\mathcal{T}}|$, namely $(x_t, z_t^*)_{t \in \mathcal{O}_{\mathcal{T}}}$, where

$$z_t^* = \begin{cases} x_t^* - x_{t^-}^* & \text{if } ht_{\mathcal{T}}(t) > 0 \\ x_t^* & \text{if } ht_{\mathcal{T}}(t) = 0 \end{cases}, \quad t \in \mathcal{O}_{\mathcal{T}}.$$

(For background on biorthogonal systems in Banach spaces we refer the reader to the book Hájek et al. (2008).) The following proposition is now immediate.

Proposition 15 – *Let Z be a Banach space not admitting an uncountable biorthogonal system and let (\mathcal{T}, \leq) be a tree such that $\mathcal{O}_{\mathcal{T}}$ is uncountable. Then $\Sigma_{\mathcal{T}}$ does not factor through Z .*

It is consistent with ZFC that there exists an Asplund space W with $Sz(W) > \omega_1$ and W does not admit an uncountable biorthogonal system. Thus, by Proposition 15, it is consistent with ZFC that the operator $\Sigma_{\mathcal{S}}$ defined earlier in the current section is not universal for $\mathcal{D} \cap \bigcap \mathcal{SZ}_{\omega_1}$. An example of a compact Hausdorff space K such that $C(K)$ is such a space W was constructed in the 1970s by Kunen, though the construction was not published until much later by Negreponitis⁴². (For further historical remarks concerning the existence of uncountable biorthogonal systems, see Remark 4 of Todorčević (2006).) Since a Banach space $C(L)$ is Asplund if and only if L is scattered⁴³, the space $C(K)$ arising from Kunen's construction is Asplund. Moreover, the Cantor-Bendixson rank of Kunen's space K is larger than ω_1 . Thus, the $C(K)$ space arising from Kunen's construction is indeed an example of such

⁴²Negreponitis, 1984, "Banach spaces and topology".

⁴³Namioka and Phelps, 1975, "Banach spaces which are Asplund spaces".

a space W once we have observed the following fact: for L a compact Hausdorff space the Szlenk index of $C(L)$ is bounded below by the Cantor-Bendixson rank of L . This is an easy consequence of the well-known fact that the mapping that takes $l \in L$ to the evaluation functional of $C(L)$ at l is a homeomorphic embedding with respect to the weak* topology, and the image of L under this embedding is a 1-separated subset of $B_{C(L)^*}$. From this fact it is easy to see that $Sz(C(L), 1)$ is bounded below by the Cantor-Bendixson rank of L , hence $Sz(C(L))$ is bounded below by the Cantor-Bendixson rank of L .

Finally, we mention a more recent construction of Brech and Koszmider⁴⁴, who establish the consistency of a scattered compact Hausdorff space J having Cantor-Bendixson rank equal to $\omega_2 + 1$ and such that $C(J)$ does not admit an uncountable biorthogonal system. If \mathcal{T} is a tree that is blossomed in the generalised sense introduced at the beginning of the current section, and if $\rho(\mathcal{T}) = \omega_2 + 1$, then $Sz(\Sigma_{\mathcal{T}}) = \omega_2 \omega$. However, by Proposition 15, $\Sigma_{\mathcal{T}}$ does not factor through the Brech-Koszmider space $C(J)$ which satisfies $Sz(C(J)) \geq \omega_2 \omega$.

Acknowledgments

The author is grateful to Ryan Causey for pointing out the relevance of the references Alspach, Judd, and Odell (2005) and Beanland and Causey (2018) to the current paper. The author also thanks the anonymous referee(s) for numerous helpful comments that have significantly improved the quality of the paper, including in particular for suggesting a significant simplification to the proof of Lemma 3.

References

- Albiac, F. and N. J. Kalton (2016). *Topics in Banach space theory*. Graduate Texts in Mathematics, No. 233 (2nd revised and updated edition). Springer International Publishing (cit. on pp. 163, 170, 187, 192, 199).
- Alspach, D. E., R. Judd, and E. Odell (2005). “The Szlenk index and local ℓ_1 -indices”. *Positivity* **9**, pp. 1–44 (cit. on pp. 166, 206).
- Beanland, K. and R. M. Causey (2017). “Genericity and universality for operator ideals”. *Preprint*. URL: <https://arxiv.org/abs/1711.09244> (cit. on p. 168).
- Beanland, K. and R. M. Causey (2018). “Quantitative factorization of weakly compact, Rosenthal, and ξ -Banach-Saks operators”. *Math. Scand.* **123** (2), pp. 297–319 (cit. on pp. 180, 206).
- Bourgain, J. (1979). “The Szlenk index and operators on $C(K)$ -spaces”. *Bull. Soc. Math. Belg. Sér. B* **31** (1), pp. 87–117 (cit. on pp. 175, 203).

⁴⁴Brech and Koszmider, 2011, “Thin-very tall compact scattered spaces which are hereditarily separable”.

References

- Brackebusch, R. E. (1988). "James Space on General Trees". *J. Funct. Anal.* **79** (12), pp. 446–475 (cit. on p. 179).
- Brech, C. and P. Koszmider (2011). "Thin-very tall compact scattered spaces which are hereditarily separable". *Trans. Amer. Math. Soc.* **363** (1), pp. 501–519 (cit. on p. 206).
- Brooker, P. A. H. (2012). "Asplund operators and the Szlenk index". *J. Operator Theory* **68**, pp. 405–442 (cit. on pp. 170–172, 197).
- Brooker, P. A. H. (2013). "Szlenk and w^* -dentability indices of the Banach spaces $C([0, \alpha])$ ". *J. Math. Anal. Appl.* **399** (2), pp. 559–564 (cit. on p. 196).
- Brooker, P. A. H. (2017). "Non-Asplund Banach spaces and operators". *J. Funct. Anal.* **273** (12), pp. 3831–3858 (cit. on pp. 168, 197).
- Causey, R. M. (2017). "The Szlenk index of injective tensor products and convex hulls". *J. Funct. Anal.* **272** (8), pp. 3375–3409 (cit. on p. 172).
- Cilia, R. and J. M. Gutiérrez (2015). "Nonexistence of certain universal polynomials between Banach spaces". *J. Math. Anal. Appl.* **427** (2), pp. 962–976 (cit. on p. 168).
- Davis, W. J. and W. B. Johnson (1973). "A renorming of nonreflexive Banach spaces". *Proc. Amer. Math. Soc.* **37**, pp. 486–488 (cit. on p. 182).
- Diestel, J., H. Jarchow, and A. Pietsch (2001). "Operator ideals". In: *Handbook of the geometry of Banach spaces, Vol. I*. Amsterdam: North-Holland, pp. 437–496 (cit. on p. 168).
- Dilworth, S. J. (1985). "Universal noncompact operators between super-reflexive Banach spaces and the existence of a complemented copy of Hilbert space". *Israel J. Math.* **52** (1–2), pp. 15–27 (cit. on p. 168).
- Dilworth, S. J. et al. (2017). "Equivalent norms with the property (β) of Rolewicz". *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas* **111** (1), pp. 101–113 (cit. on pp. 164, 165, 181).
- Fabian, M. et al. (2001). *Functional analysis and infinite-dimensional geometry*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 8. New York: Springer-Verlag (cit. on p. 192).
- Gasparis, I. (2005). "Operators on $C(K)$ spaces preserving copies of Schreier spaces". *Trans. Amer. Math. Soc.* **357** (1), pp. 1–30 (cit. on pp. 174–176).
- Girardi, M. and W. B. Johnson (1997). "Universal non-completely-continuous operators". *Israel J. Math.* **99**, pp. 207–219 (cit. on p. 168).
- Godefroy, G., N. J. Kalton, and G. Lancien (2001). "Szlenk indices and uniform homeomorphisms". *Trans. Amer. Math. Soc.* **353** (10), pp. 3895–3918 (cit. on p. 173).
- Grothendieck, A. (1953). "Sur les applications linéaires faiblement compactes d'espaces du type $C(K)$ ". *Canadian J. Math.* **5**, pp. 129–173 (cit. on p. 199).
- Hagler, J. and E. Odell (1978). "A Banach space not containing ℓ_1 whose dual ball is not weak* sequentially compact". *Illinois J. Math.* **22** (2), pp. 290–294 (cit. on p. 179).

- Hájek, P. et al. (2008). *Biorthogonal systems in Banach spaces*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 26. New York: Springer (cit. on p. 205).
- Hinrichs, A., E. Novak, and H. Woźniakowski (2013). “Discontinuous information in the worst case and randomized settings”. *Math. Nachr.* **286** (7), pp. 679–690 (cit. on p. 168).
- Hinrichs, A. and A. Pietsch (2000). “The closed ideal of (c_0, p, q) -summing operators”. *Integral Equations and Operator Theory* **38** (3), pp. 302–316 (cit. on p. 168).
- Johnson, W. B. (1971). “A universal non-compact operator”. *Colloq. Math.* **23**, pp. 267–268 (cit. on pp. 167, 198, 203).
- Johnson, W. B. and H. P. Rosenthal (1972). “On w^* -basic sequences and their applications to the study of Banach spaces”. *Studia Math.* **43**, pp. 77–92 (cit. on pp. 164, 166, 187, 189–192).
- Kadec, M. Ī. (1958). “On strong and weak convergence”. Russian. *Dokl. Akad. Nauk SSSR* **122**, pp. 10–13 (cit. on p. 182).
- Klee, V. (1960). “Mappings into normed linear spaces”. *Fund. Math.* **49**, pp. 25–34 (cit. on p. 182).
- Lancien, G. (1996). “On the Szlenk index and the weak*-dentability index”. *Quart. J. Math. Oxford Ser. (2)* **47** (185), pp. 59–71 (cit. on pp. 164–166, 181, 187, 194).
- Lancien, G. (2006). “A survey on the Szlenk index and some of its applications”. *RACSAM Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat.* **100**, pp. 209–235 (cit. on pp. 163, 171).
- Lindenstrauss, J. and A. Pełczyński (1968). “Absolutely summing operators in \mathcal{L}_p -spaces and their applications”. *Stud. Math.* **29**, pp. 275–326 (cit. on pp. 167, 198, 203).
- Lindenstrauss, J. and L. Tzafriri (1977). *Classical Banach spaces. I. Sequence spaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Vol. 92. Berlin: Springer-Verlag (cit. on pp. 182, 197).
- Namioka, I. and R. R. Phelps (1975). “Banach spaces which are Asplund spaces”. *Duke Math. J.* (4), pp. 735–750 (cit. on p. 205).
- Negrepontis, S. (1984). “Banach spaces and topology”. In: *Handbook of set-theoretic topology*. Amsterdam: North-Holland, pp. 1045–1142 (cit. on p. 205).
- Nyikos, P. J. (1997). “Various topologies on trees”. In: *Proceedings of the Tennessee Topology Conference (Nashville, TN, 1996)*. River Edge, NJ: World Sci. Publ., pp. 167–198 (cit. on p. 176).
- Oikhberg, T. (2016). “A note on universal operators”. In: *Ordered structures and applications. Positivity VII (Zaanen centennial conference), Leiden, the Netherlands, July 22–26, 2013*. Basel: Birkhäuser/Springer, pp. 339–347 (cit. on p. 168).
- Pietsch, A. (1980). *Operator ideals*. North-Holland Mathematical Library, Vol. 20. Amsterdam: North-Holland Publishing Co. (cit. on pp. 167, 170).
- Rosenthal, H. P. (2003). “The Banach spaces $C(K)$ ”. In: *Handbook of the geometry of Banach spaces, Vol. 2*. Amsterdam: North-Holland, pp. 1547–1602 (cit. on p. 163).

References

- Samuel, C. (1984). "Indice de Szlenk des $C(K)$ (K espace topologique compact dénombrable)". In: *Seminar on the geometry of Banach spaces, Vol. I, II (Paris, 1983)*. Vol. 18. Publ. Math. Univ. Paris VII. Paris: Univ. Paris VII, pp. 81–91 (cit. on p. 197).
- Sersouri, A. (1989). "Lavrientiev index for Banach spaces". *C. R. Acad. Sci. Paris Sér. I Math.* **309** (2), pp. 95–99 (cit. on p. 172).
- Singer, I. (1970). *Bases in Banach spaces. I*. Die Grundlehren der mathematischen Wissenschaften, Band 154. New York: Springer-Verlag (cit. on p. 192).
- Stegall, C. (1975). "The Radon-Nikodým property in conjugate Banach spaces". *Trans. Amer. Math. Soc.* **206**, pp. 213–223 (cit. on p. 167).
- Stegall, C. (1981). "The Radon-Nikodým property in conjugate Banach spaces. II". *Trans. Amer. Math. Soc.* **264** (2), pp. 507–519 (cit. on p. 167).
- Szlenk, W. (1968). "The non-existence of a separable reflexive Banach space universal for all separable reflexive Banach spaces". *Studia Math.* **30**, pp. 53–61 (cit. on p. 163).
- Todorčević, S. (1984). "Trees and linearly ordered sets". In: *Handbook of set-theoretic topology*. Amsterdam: North-Holland, pp. 235–293 (cit. on p. 172).
- Todorčević, S. (2006). "Biorthogonal systems and quotient spaces via Baire category methods". *Math. Ann.* **335** (3), pp. 687–715 (cit. on p. 205).

Contents

1	Introduction	163
1.1	Structure of Banach spaces with large Szlenk index	164
1.2	Universal operators	166
1.3	Notation and terminology	169
2	The Szlenk index, trees and operators on Banach spaces over trees	171
2.1	The Szlenk index	171
2.2	Trees	172
2.3	Operators on Banach spaces over trees	177
3	Absolutely convex sets of large Szlenk index	180
4	Structure of Banach spaces of large Szlenk index	192
5	Universal operators of large Szlenk index	195
6	When the codomain is separable	198
7	Uncountable Szlenk indices and universality	204
	Acknowledgments	206
	References	206
	Contents	i